

Akaniaceae, a new family record for Laos

SOULIVANH LANORSAVANH^{1,*}, VICHITH LAMXAY¹, SAWAI MATTAPHA²,
KEOUDONE SOUVANNAKHOUUMANE¹, KHAMFA CHANTHAVONGSA¹ & SOMCHANH BOUNPHANMY¹

ABSTRACT

A collection from Phou Chom Voy Provincial Protected Area, Bolikhamsai, represents the first records for the family Akaniaceae as well as the genus *Bretschneidera* and the species *Bretschneidera sinensis* in Laos. Here we report the new and extended distributional range. A detailed description and photographs are provided for easy identification.

KEYWORDS: Brassicales, Bretschneideraceae, extended distribution, Indo-China.

Accepted for publication: 11 October 2021. Published online: 25 October 2021

INTRODUCTION

The high moist mountainous area “Phou Chom Voy”, part of the Annamite Mountain Range, is located in the Phou Chom Voy Provincial Protected Area of Bolikhamsay Province. It extends to the Laotian-Vietnamese border and encompasses the Khamkeut and Xaychamphone Districts. “Phou Chom Voy” has been shown to have a rich biodiversity and natural water resources, as well as potential recreational and tourist opportunities. It is a diversity hotspot of taxa, home to moist and clouded high montane evergreen forests, at elevations of 800–1,820 m asl. The area is difficult to access and probably contains many endemic and unexplored species. At present, scientific data of the flora from this area are insufficient (V. Lamxay pers. comm.).

In the past few years, there were two new records of orchids for Laos, namely *Odontochilus elwesii* C.B.Clarke ex Hook.f. and *O. putaoensis* X.H.Jin, L.A.Ye & A.T.Mu. (Lanorsavanh *et al.*, 2019). A year later, the new species, *Begonia phouchomvoyensis* Lanors., Lamxay & Souvann., was discovered from the same area (Lanorsavanh *et al.*, 2020).

Botanical surveys in the Phou Chom Voy Provincial Protected Area were undertaken between 2018 to 2021. Some plant collections, identified with

Hemsley (1901), Niyomdharm (1991), Lu & Boufford (2005) and Gardner *et al.* (2007), appeared to be *Bretschneidera sinensis* Hemsl. (Akaniaceae), a new family, genus and species record for Laos. Vouchers are deposited in the Biology Herbarium of the National University of Laos and distributed to the National Herbarium of Laos (HNL).

Bretschneidera Hemsl. was formerly placed in the family Sapindaceae by Hemsley (1901) and later in Bretschneideraceae with a possible close relationship to Hippocastanaceae and Sapindaceae (Boufford *et al.*, 1989; Tobe & Peng, 1990; De Craene *et al.*, 2002; Lu & Boufford, 2005). More recently, proof was provided that the genus has to be included in the family Akaniaceae Stapf (Soltis *et al.*, 2000; APG IV, 2016). This position was already earlier pointed out by Tobe & Peng (1990), Rodman (1991), Rodman *et al.* (1993), Tobe & Raven (1995), Doweld (1996) and De Craene *et al.* (2002).

Akaniaceae, a small family of flowering plants classified in the order Brassicales, comprises two monotypic genera, *Akania* Hook.f., a native species of Australia, and *Bretschneidera*, confined to South-East Asia (Lu & Boufford, 2005; APG IV, 2016). The family is characterized by combinations of being a tree, having compound leaves, opposite to

¹ Department of Biology, Faculty of Natural Science, National University of Laos, Vientiane, Laos.

² Department of Biology, Faculty of Science, Udon Thani Rajabhat University, Udon Thani 41000, Thailand.

* Corresponding author: s.lanorsavanh@nuol.edu.la

subopposite leaflets, absence of stipels, bisexual flowers, campanulate calyces with 5 lobes, corolla with 5 lobes, a superior ovary with 3–5 locules, axile placentation and capsular fruits.

The genus *Bretschneidera* differs from *Akania* by having imparipinnate leaves (vs paripinnate in *Akania*), entire leaflet margins (vs serrate in *Akania*), a racemose inflorescence (vs paniculate inflorescence in *Akania*), zygomorphic flowers (vs actinomorphic flowers in *Akania*) and presence of a floral disk (vs absent in *Akania*).

In this communication, we report *Bretschneidera*, a monotypic genus, and *Bretschneidera sinensis* as new records for Laos. Previously, the species was reported ranging from India, southern China and northern Vietnam to confined areas in Northern Thailand (Doi Phu Kha National Park, Nan Province; Niyomdharm, 1991; Gardner *et al.*, 2007; Kumar *et al.*, 2017), but now it is known to have a wider distribution, extending into Laos. The species was not yet recorded in the plant check-list of Laos (Newman *et al.*, 2007; Jin *et al.*, 2016), and the online check-list, which is continuously updated by the Royal Botanic Garden Edinburgh (<https://padme.rbge.org.uk/laos/list/>). Although, it was assumed to occur in Laos by Lu & Boufford (2005); Wang *et al.* (2018), but no specimen was recorded yet.

NEW RECORD

Akaniaceae Stapf, Bull. Misc. Inform. Kew, 1912: 380. 1912; C.Bayer & O.Appel in Kubitzki, Fam. & Gen. Vasc. Pl. 5: 21. 2002; APG IV, Bot. J. Linn. Soc. 181: 18. 2016; Chamch. *et al.*, Threat. Pl. Thailand. 25. 2017. Type: *Akania* Hook f.

Trees. Leaves alternate, paripinnate or imparipinnate; stipules present or absent; veins pinnate; stipels absent. Inflorescence racemose or paniculate, axillary or terminal, erect; bracts present. Flowers bisexual, actinomorphic or zygomorphic, 5-merous, white or pink. Calyx broadly campanulate, 5-lobed. Petals 5, convolute or imbricate. Stamens 5–10; filaments filiform, hairy; anthers dorsifixed, opening length-wise. Disk absent or present. Ovary superior, 3–5 locules; placentation axile; ovules 2 per locule, pendulous; style longer than stamens; stigma capitate, small. Fruit a capsule, ellipsoid-globose to subglobose

to ovoid or obovoid to pyriform; loculicidally dehiscent. Seeds red or yellow, ellipsoid-globose to ovoid, compressed, without endosperm and aril.

Two genera: *Akania* and *Bretschneidera*. The first genus is native to Australia. Only *Bretschneidera* was previously recorded from India, S China, Taiwan, N Thailand and N Vietnam. It is now also known for Laos.

Bretschneidera Hemsl., Hooker's Icon. Pl. 28(1): 2708. 1891; L.L.Lu & Boufford in C.Y.Wu & P.H.Raven, Fl. China 8: 197. 2005; C.Bayer & O.Appel in Kubitzki, Fam. & Gen. Vasc. Pl. 5: 23. 2002; Chamch. *et al.*, Threat. Pl. Thailand: 25. 2017. Type: *Bretschneidera sinensis* Hemsl.

Tree 10–25 m tall. Bark grey-brown, young twigs with lenticels and leaf scars. Leaves imparipinnate, with 8–31 leaflets; venation pinnate; stipules present, caducous; leaflets opposite or lower ones alternate, margin entire. Inflorescence racemose, terminal. Flowers zygomorphic. Calyx 5-lobed. Petals 5, imbricate, free, adnate at middle to upper part of calyx tube, upper petal hood-like, covering stamens and style. Stamens 8 (4+4). Disk present. Ovary 3-locular, hairy; ovules pendulous; style apically bent abruptly downward.

A monospecific genus; *Bretschneidera sinensis*; previously known from India, S China, Taiwan, N Thailand and N Vietnam, it is now also known for Laos.

Bretschneidera sinensis Hemsl., Hooker's Icon. Pl. 28: t. 2708. 1901; Gagnep. in Lecomte, Fl. Indo-Chine, Suppl. 1: 1001. 1950; Santisuk, Nat. Hist. Bull. Siam Soc. 37(2): 173. 1989; Niyomdharm in Smitinand & K.Larsen, Fl. Thailand 5(3): 239. 1991; L.L.Lu & Boufford in C.Y.Wu & P.H.Raven, Fl. China 8: 197. 2005. Type: China, Szemao, Yunnan, at 5,000 ft., A. Henry 11651 (lectotype K [K000681003!], designated by Kumar *et al.* 2017; isolectotypes A [A00050653!], K [K000681002!]). Fig. 1.

— *Bretschneidera yunshanensis* Chun & F.C.How, Acta Phytotax. Sin. 7(1): 68. 1958. Type: China, Hunan, Hengshan, Fangkwang, Tzu, 9 May 1955, K.H. Chu 99 (holotype SCBI).




Figure 1. *Bretschneidera sinensis* Hemsl.: A. Leafy twig with inflorescences; B. Bark; C. Flowers; D. Corolla with 5 lobes (dissected). E. Calyx (corolla lobes removed) showing inside of calyx, attachment of ovary and stamens (above) and outside of calyx with minute lobes (below). F. Stamens. Scale bars: 1 cm. All from SL2157 by S. Lanorsavanh.

Leaves: petioles 11–17.2 cm long; rachis 11–21 cm long, pubescent; ultrajugal part 1.4–4.2 cm long, slightly greenish, reddish adaxially, sparsely hairy, glabrescent. Leaflets 7–17; petiolules 3–9 mm long; lamina obliquely lanceolate, oblong-lanceolate, ovate-lanceolate or elliptic, 6–19.5 × 2.7–7.7 cm, chartaceous, base obtuse to rounded to cuneate, apex

acute to acute or caudate, adaxial surface green to somewhat pale green, glabrous, abaxial surface grey, minutely pubescent, lateral veins 10–16 pairs. **Racemes** 12–30 cm long, hairy. **Flowers** pink or pink-white or rose-white. **Pedicels** 2–3.2 cm long, hairy. **Calyx** broadly campanulate, 1.1–1.7 × 2–2.5 cm; lobes broadly triangular, 2–3 mm long, hairy

on both sides. *Petals* broadly spatulate to obovate, 1.8–2.7 × 1.5–2.2 cm, base cuneate, apex weakly notched; upper petal pink to white, becoming darker with age, red striate. *Stamens*: filaments 2.2–2.5 cm long, hairy at base, curving downward apically; anthers oblong, 5–6.5 × 1.5–2 mm. *Disk* annular. *Ovary* ellipsoid-globose to subglobose to ovoid or obovoid, 3–4 × ca 3 cm; style 2.2–2.4 cm long, hairy; stigma ca 1 mm diam. *Fruit* not seen.

Distribution.— See under genus.

Ecology & Phenology.— In Laos collected near a stream in evergreen forest at ca 1,100 m alt. Flowering in March.

Vernacular.— ດອກພວງບົວຈອມວອຍ (Dok phuang boua chom voy).

Specimens examined.— Laos: Bolikhamsai, Khaneut District, Nahueng Village, Phou Chom Voy Provincial Protected Area, 18°25'10.3"N, 105°03'13.4"E, 1,084 m, 3 Mar. 2021, *Lanorsavanh et al. SL2157 (HNL, NLS*=National University of Laos, Faculty of Natural Science).

Conservation status— The species is present with less than a 100 individual trees in Laos over an area of ca 10 km² that is only at one site a Protected Area; therefore, in Laos it is rare and Endangered (Sun, 1998).

Note.— *Bretschneidera sinensis* is characterized by its imparipinnate leaves, erect racemes, 5-lobed corolla, corolla lobes broadly spatulate to obovate, base cuneate, upper petal hood-like covering the stamens and the style, 8 stamens and a 3-locular ovary.

ACKNOWLEDGEMENTS

The authors would like to thank the staff of the Department of Biology, Faculty of Natural Science, National University of Laos and the Department of Biology, Faculty of Science, Udon Thani Rajabhat University, Thailand, for use of their facilities during the study and the staff of the herbaria for their collaboration. We are especially grateful to the Environment Protection Fund (EPF) under the Project of Capacity Enhancement of Laos's Natural Sciences, in the Sub-Project Number PICE-LENS2-010, for supporting the field survey at Phou Chom Voy Provincial Protected Area. We also express

our gratitude to Pongkham Phommachan for assisting during the field work. Finally, thanks to Jeffrey Nash who kindly reviewed the English and two reviewers for helpful comments for improving the manuscript.

REFERENCES

APG IV (2016). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181 (1): 1–20.

Boufford, D.E., Kjaer, A., Madsen, J.O. & Skrydstrup, T. (1989). Glucosinolates in Bretschneideraceae. *Biochemical Systematics and Ecology* 17: 375–379.

De Craene, L.P.R., Yang, T.Y.A., Schols, P. & Smets, E.F. (2002). Floral anatomy and systematics of *Bretschneidera* (Bretschneideraceae). *Botanical Journal of the Linnean Society* 139: 29–45.

Doweld, B.A. (1996). The Carpology and Taxonomic relationships of *Bretschneidera* (Bretschneideraceae). *Acta Botanica Malacitana* 21: 79–90.

Gardner, S., Sirisunthorn, P. & Anusarnsunthorn, V. (2007). A field guide to Forest Trees of Northern Thailand: 139. Kobfai publishing project, Bangkok. p 139.

Hemsley, W.B. (1901). *Bretschneidera sinensis*. Hooker's *Icones Plantarum* 28: t. 2708.

Jin, H.Y., Ahn, T.H., Lee, H.J., Song, J.H., Lee, C.H., Kim, Y.J., Yoon, J.W., Chamg, K.S., Kang, H.S., Cheng, H.C., Park, J.H., Bang, M., Bouahom, B., Phongoudome, C., Thammavong, V., Soukladeth, P., Bouamanivong, S., Phasaysombath, T., Chanthavongsa, K., Lanorsavanh, S., Sayyalath, S., Bounithiphonh, C. & Phonephanome, P. (2016). A Checklist of Plants in Lao PDR. Korea National Arboretum of the Korea Forest Service.

Kumar, V., Dash, S.S., Panday, S., Lahiri, S., Sinha, B.K. & Sihgh, P. (2017). Akanaceae: a new family record for flora of India and lectotypification of the name *Bretschneidera sinensis*. *Nelumbo* 59(1): 1–9.

Lanorsavanh, S., Hughes, M., Souvannakhoummane, K. & Lamxay, V. (2020). *Begonia phouchomvogensis* (Begoniaceae), a new species from Laos. *Taiwania* 65(1): 37–40.

Lanorsavanh, S., Souvannakhouummane, K., Kumar, P., Gale, S.W., Lamxay, V., Bounphanomy, S. & Chanthavongsa, K. (2019). Two new distribution records of *Odontochilus* Blume (Orchidaceae) from Laos. *Thai Journal of Botany* 11 (2): 85–93.

Lu, L.L. & Boufford, D.E. (2005). Bretschneideraceae. In: C.Y. Wu & P.H. Raven (eds), *Flora of China* 8: 197. Science Press, Beijing; Missouri Botanical Garden, St. Louis.

Newman, M., Ketphanh, S., Svengsuksa, B., Thomas, P., Sengdala, K., Lamxay, V. & Armstrong, K. (2007). A checklist of the vascular plants of Lao PDR. Royal Botanic Garden, Edinburgh.

Niyomdham, C. (1991). Bretschneideraceae. In: T. Smitinand & K. Larsen (eds), *Flora of Thailand* 5: 239–240. The Forest Herbarium, Bangkok.

Rodman, J.E. (1991). A taxonomic analysis of glucosinolate-producing plants. Part 1. Phenetics. *Systematic Botany* 16: 589–618.

Rodman, J.E., Price, R.A., Karol, K., Conti, E., Sytsma, K.J. & Palmer, J.D. (1993). Nucleotide sequences of the *rbcL* gene indicate monophyly of mustard oil plants. *Annals of the Missouri Botanical Garden* 80: 686–699.

Soltis, D.E., Soltis, P.S., Chase, M.W., Mort, M.E., Albach, K.C., Zanis, M., Savolainen, V., Hanh, W.H., Hoot, S.B., Fay, M.F., Axtell, M., Swensen, S.M., Prince, L.M., Kress, W.J., Nixon, K.C. & Farris, J. (2000). Angiosperm phylogeny inferred from 18S rDNA, *rbcL*, and *atpB* sequences. *Botanical Journal of the Linnean Society* 133: 381–461.

Sun, W. (1998). *Bretschneidera sinensis*. The IUCN Red List of Threatened Species 1998: e.T32324A9697750. Downloaded on 08 October 2021.

Tobe, H. & Peng C.-I (1990). The embryology and taxonomic relationships of *Bretschneidera* (Bretschneideraceae). *Botanical Journal of the Linnean Society* 103(2): 139–152.

Tobe, H. & Raven, P.H. 1995. Embryology and relationships of *Akania* (Akaniaceae). *Botanical Journal of the Linnean Society* 118: 261–274.

Wang, M.N., Duan, L., Qiao, Q., Wang, Z.F., Zimmer, E.A., Li, Z.C. & Chen, H.F. (2018). Phylogeography and conservation genetics of the rare and relict *Bretschneidera sinensis* (Akaniaceae) PLOS ONE 13(1): e0189034.