



Original article

Using *in situ* seed baiting technique to isolate and identify endophytic and mycorrhizal fungi from seeds of a threatened epiphytic orchid, *Dendrobium friedericksianum* Rchb.f. (Orchidaceae)



Na-monrug Khamchatra,^{a, e} Kingsley Dixon,^b Kongkanda Chayamarit,^c
Somsak Apisitwanich,^d Sureeya Tantiwiwat^{a, *}

^a Department of Botany, Faculty of Science, Kasetsart University, Bangkok, Thailand

^b Kings Park and Botanic Garden, Perth Western, Australia

^c The Forest Herbarium, Department of National Park, Wildlife and Plant Conservation, Bangkok, Thailand

^d Department of Genetics, Faculty of Science, Kasetsart University, Bangkok, Thailand

^e Department of Biology, Faculty of Science and Technology, Rambhai Barni Rajabhat University, Chanthaburi, Thailand

ARTICLE INFO

Article history:

Received 16 June 2015

Accepted 20 August 2015

Available online 14 January 2016

Keywords:

Dendrobium friedericksianum

Endophytic fungi

In situ

Mycorrhizal fungi

Tulasnella

ABSTRACT

All orchids require association with mycorrhizal fungi for seed germination and development under natural conditions but their identification and feasibility are not known. The *in situ* seed baiting germination was examined of *Dendrobium friedericksianum* Rchb.f., a native threatened epiphytic orchid species, to detect fungi that promote germination. It was found that seed germination percentages were lowered by 0.1%, with a total of seven protocorms formed. Six endophytic fungi were isolated from seven protocorms in seed packets. Three binucleate *Rhizoctonia*-like fungal isolates which formed a teleomorphic state were morphologically identified as *Tulasnella violea*, *Epulorhiza repen* (anamorph *Tulasnella*) and *Trichosporiella multisporum*. The species of *Beauveria* and *Fusarium* which are endophyte fungi were also isolated and may play an important role for plant growth and survival of *D. friedericksianum*. The results in this study suggested that *in situ* seed baiting is beneficial for screening compatible mycorrhizal fungi to promote the growth and propagation of epiphytic orchids.

Copyright © 2016, Kasetsart University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Dendrobium is one of the largest genera of Orchidaceae, with over 1000 species worldwide (Leitch et al., 2009). Many *Dendrobium* orchids are important horticultural crops and have been used in commercial trade due to their popular flowering abundance, wide range of flower colors, sizes, shapes, year-round availability and lengthy post-harvest life (Hossain, 2011). Most *Dendrobium* species are suspected of becoming extinct. Of these, *Dendrobium friedericksianum* Rchb.f. is a native epiphytic orchid found in tropical rain forests in Eastern Thailand according to the checklist of international species, (Convention on International Trade in Endangered Species of Wild Fauna and Flora, 2014). Due to their beautiful, bright golden-yellow flowers with diameters

around 4–5 cm each, the orchid has attracted collectors and so specimens are in high demand for illegal trading. Consequently, this species is on the verge of being seriously endangered in the wild.

Orchid seeds are usually small, dust-like and lacking an endosperm. Consequently, germination of orchid seeds and seedling development need compatible endophytic mycorrhizal fungi to provide the minute seeds with carbon, water and nutrients under natural conditions (Arditti and Ghani, 2000; Rasmussen and Rasmussen, 2009). Asymbiotic seed germination techniques and *in vitro* rapid propagation of seedlings have been applied to *D. friedericksianum* (Prasertsivivatna and Koolpluksee, 2011), but the seedlings have rarely survived after a transplant to nature despite being well grown in aseptic culture. It is assumed that they may be dependent on suitable mycorrhizal fungi for orchid seedling survival.

Most studies on orchid mycorrhizal fungi have been focused on *in vitro* symbiotic techniques using fungal isolates obtained from

* Corresponding author.

E-mail addresses: namonrug.k@gmail.com (N.-m. Khamchatra), sureeya50@hotmail.com (S. Tantiwiwat).

adult orchid roots, not from protocorms or seedling stages (Chutima et al., 2011; Nontachaiyapoom et al., 2011). It was not clear whether mycorrhizal fungi found in adult orchid roots are the same fungi necessary for seed germination until Rasmussen and Whigham (1993) studied and developed *in situ* seed germination methods which involved burying seed packets in the field during the growing season to assess the presence of fungi capable of supporting germination. This seed baiting has been successfully used by McKendrick et al. (2000) and Brundrett et al. (2003).

The present study hypothesized that the species-specific mycorrhizae isolated from the developed protocorms should have a greater impact on the success of germination and subsequent stages of seed development. The aim of this study was to isolate and identify mycorrhizal and non-mycorrhizal endophytic fungi from protocorms of *D. friedericksianum* using the *in situ* seed baiting method. This study is the first to report on culturable mycorrhizal fungi that may be valuable for orchid propagation and conservation.

Materials and methods

Study site and seed collection

In December 2012, 9–10 mth-old, hand-pollinated, yellow-green capsules of *D. friedericksianum* were collected from fruit orchards in Chanthaburi, a southeastern province of Thailand. The capsules were put in a paper bag and transported to Rambhai Barni Rajabhat University. The capsules were kept in silica gel-desiccated plastics boxes in the dark for approximately 1 wk at room temperature. The seeds were then transferred to sealed glass bottles stored at 4 °C in the dark until use, within 4–6 mth (Stewart and Kane, 2007).

In situ baiting method

In January 2013, approximately 300–400 *Dendrobium friedericksianum* seeds were placed in separate folding 5 cm × 10 cm nylon mesh pouches with a 45 µm pore size, which was large enough to allow fungal hyphae to enter but small enough to retain seeds inside the pouch. Each of these packets was placed inside a 35 mm plastic slide mount (Brundrett et al., 2003; Zi et al., 2014). All packets were secured onto the tree bark near the adult roots of *D. friedericksianum* and were then covered with a thin layer of sterile wet *Sphagnum* moss to prevent desiccation but suitable for light to penetrate. In July 2013, harvests were taken during the orchid growing season. Recovered seed packets were placed between moist sheets of sterilized wet cotton wool and stored at 4 °C.

Seed germination and development assessment

Seed germination and development were observed under a dissecting stereomicroscope and scored on a scale of 0–5 as described by Arditti and Ghani (2000). Seed germination and protocorm development percentages were calculated.

Fungal isolation from seed baiting packets

Mycorrhizal fungi were isolated from the collected germinated seeds which had approached stage 3 according to Brundrett et al. (2003) with slight modifications before dispersal on the agar surface as follows. The seed packet surfaces were washed with running tap water and sterilized by immersing in 70% ethanol for 15–30 s depending on the seed size. The seed packets were then dipped in sodium hypochlorite solution containing 0.5% available chlorine for 3–5 min and rinsed three times with sterile double-distilled water. They were opened under a dissecting microscope. The protocorms

were then sown on modified Czapek Dox agar medium (Yamato et al., 2005) supplemented with 0.05 g/L streptomycin. The pH was adjusted to 7.3. After 4–7 d, the fungal hyphae growth that had emerged from the protocorms were excised and transferred onto fresh potato dextrose agar (PDA) and incubated at 25 ± 2.0 °C in the dark. Fungal colonies from actively growing isolates were subcultivated 2–3 times onto a fresh PDA dish to be purified and assigned a strain number (denoted with the code DFCT-B).

Cultural and morphological characterization

The growth rate and colony characteristics were recorded from the culture grown on PDA (Sneh et al., 1991; Currah et al. 1997; Roberts, 1999). Microscopic observations were made using a slide culture technique. Fungal hyphae stained with lactophenol cotton blue and moniloid cells were assessed visually and microscopically using light microscopy (Eclipse E400; Nikon; Tokyo, Japan). The number of nuclei per cell was determined after staining with safranin O–KOH using Bandoni's method (Bandoni, 1979).

Results

In situ symbiotic seed germination

The 30 seed packets placed *in situ* on the bark of trees in January 2013 (Fig. 1) were collected 7 mth later in July 2013. The numbers of seeds that reached or passed each germination stage out of the total seeds placed in each packet as successful events for the specific stage were counted. The results of seed germination are shown in Table 1. The total seed germination percentage of *D. friedericksianum* was 77.8 ± 9.3. Seed germination of orchid species up to the total achievement of stage-3 protocorms (appearance of pro-meristems) were determined (Fig. 2A). The peloton and coiled mycelium (Fig. 2B) of the orchid's mycorrhizal fungi were observed in the protocorms. More than 50% of germinated seeds developed to stage-2.

Microbiont identification and morphological characteristics

In total, six endophyte mycorrhizal fungi were isolated from seven protocorms in the formation stage collected *in situ*. These isolates were divided into two groups based on cultural characteristics. Group I, containing isolates DFCT-B-05, DFCT-B-09 and DFCT-B-17 showed resemblance to *Rhizoctonia*-like orchid endophytes consisting of colorless colonies, right-angle branching, constrictions at branch point, septum in the branch hyphae near its point of origin, sclerotia formation, chains of moniloid cells and a number of nuclei in young cells. These three isolates are described below.

Isolate DFCT-B-05 grew slowly with a white-cream colony, submerged and reached 9 cm in diameter after 2 wk on PDA (Fig. 3A). Vegetative hyphae were 5–6 µm wide, binucleate, but no clamp connection was found. Basidia were swollen at the tips of subtending hyphae and bearing small, inverted-drop-shaped basidiospores, 4.5–5.5 µm (Fig. 3B) in length. This fungal isolate was identified to be *Tulasnella violea* (teleomorph).

Isolate DFCT-B-09 colony became white-to-cream in 1 wk and was submerged with glabrous surface (Fig. 3C). Hyphae were 3.5–4.0 µm wide, septate, binucleate and had a light brown, small sclerotia. Moniloid cells were ellipsoidal to spherical (Fig. 3D), hyaline were in unbranched short chains and it was identified as *Epulorhiza repens* (anamorph).

Isolate DFCT-B-17 colony was flat with white aerial hyphae (Fig. 3E). Fertile hyphae were 2.0–3.0 µm wide, septate and bearing a lateral conidia. The isolate, described as *Trichosporiella*



Fig. 1. Seed packet on a tree: (A) Seed packet covered with *Sphagnum* moss and gutter mesh; (B) Cluster of seed packets affixed to tree bark.

Table 1
Different seed germination stages of *D. friedericksianum*.

Seed germination stage	Characteristics	Percentage of seed and protocorms
0	No germination, seed with intact seed coat	22.1 ± 9.3
1	Enlarged embryo, seed coat ruptured by the enlargement of embryo (=germination)	20.0 ± 14.1
2	Continued embryo enlargement, rupture of testa	57.7 ± 13.3
3	Appearance of protomeristem	0.1 ± 0.3
4	Emergence of first leaf	0.0 ± 0.0
5	Elongation of first leaf and further development	0.0 ± 0.0

Values are mean ± standard deviation of 30 replicates.

multisporum, was globose to ellipsoid, sessile to short-pedicellate, sporulated (Fig. 3F), and featured smooth-walled conidia.

The other three isolates in Group II consisted of isolates DFCT-B-32, DFCT-B-41 and DFCT-B-47 belonging to the genus *Fusarium*. The




Fig. 2. *D. friedericksianum* protocorms: (A) Appearance of protomeristem (white arrow); (B) Peloton colonization in protocorm (black arrows).

colony's upper surface became white-creamy and the lower surface became creamy with brown zonation in 1 wk (Fig. 4A). Microconidia were ellipsoid, colorless, 2.5–4.0 µm wide and 4.0–6.0 µm in length. Macroconidia were 3.5–5.5 µm wide and 25–33 µm in length. Macroconidial morphology was slightly curved and thick, having 3–4 septa (Fig. 4B).

Isolates DFCT-B-32, DFCT-B-41 and DFCT-B-47 were saprophytic fungi. The colony of isolate DFCT-B-32 was flat and round with a white color—similar to the hyphae hyaline film from growing mycelia. It was downy in the center (Fig. 4C). Conidiogenous cells were globose to ellipsoid, smooth, short, unbranched and belonged to *Beauveria* sp. (Fig. 4D).

Discussion

D. friedericksianum is a wild epiphytic orchid found in tropical rain forests in Eastern Thailand, especially in Chanthaburi province. The unique and beautiful inflorescence of this orchid is rather expensive and popular among orchid growers. There is high demand resulting in its removal from its natural habitat for illegal trading and consequently, it is close to becoming extinct. Rajamangala University of Technology, Tawan-ok Chanthaburi campus, has conducted a research study on an asymbiotic propagation method and has released propagated orchids into natural habitats every August at both Namtok Phlio National Park and Khao Hitchakut National Park. In the survey period (2011–2012), the survival rate of *D. friedericksianum* asymbiotic seedlings was low—not considered a successful conservation and reintroduction effort. It may have been lacking mycorrhizal fungi associations to assist plant growth and fitness (Brundrett et al., 2003). The use of germinated seeds together with species-specific mycorrhizal fungi could improve the conservation programs.

To conserve and reintroduce threatened and endangered orchid species, it is necessary to propagate these species with a preference for symbiotic approaches. For this reason, symbiotic germination has become a popular and useful method for orchid seed propagation particularly where specific species are required for reintroduction programs. Up to date, the number of studies reporting symbiotic germination of endangered orchid species has been increasing (Chutima et al., 2011). The *in situ* seed baiting technique has been successfully used to detect the inoculum potential of mycorrhizal fungi and for orchid-mycorrhizal-specific investigation that supports early seed germination and protocorm development




Fig. 3. Cultural and morphological characteristics of mycorrhizal fungi on PDA: (A) *Tulasnella violea*; (B) Basidiospores of *T. violea* (white arrow); (C) *Epulorhiza* sp.; (D) Monilioid cells of *Epulorhiza* sp. (white arrow); (E) *Trichosporiella multisporum*; (F) Conidiospore of *Trichosporiella multisporum* (white arrow).

in natural habitats (Bidartondo and Read, 2008). Recently, most studies have focused on terrestrial orchid seed baiting but a few studies reported *in situ* baiting for epiphytic orchids species, *Dendrobium officinale*, *Dendrobium nobile* and *Dendrobium aphyllum* (Wang et al., 2011; Zi et al., 2014). Nevertheless, there is no report on *in situ* baiting, symbiotic seed germination of endangered wild *D. friedericksianum*. This is also the first report of protocorm formation in *D. friedericksianum* using the *in situ* baiting technique. The current results support the hypothesis that mycorrhizal fungi isolated from developed protocorms would enhance germination and seed development. The current study found a high seed germination percentage with a total of seven at protocorm stage-3. The six isolates extracted from protocorms could be divided into two groups based on morphological characteristics. Group I—three

binucleate *Rhizoctonia*-like fungal isolates which formed a teleomorphic state—were identified as *Tulasnella violea*, *Epulorhiza repen* (anamorph *Tulasnella*) and *Trichosporiella multisporum*. These fungal genera are well known as mycorrhizal fungi and are associated with a diverse range of many terrestrial and epiphytic orchids, including *Caladenia*, *Cymbidium*, *Dendrobium*, *Goodyera*, *Paphiopedilum* and *Thelymitra* (Athipunyakom et al., 2004; Nontachaiyapoom et al., 2010). Chen et al. (2012) revealed that many fungal species isolated from several species of *Dendrobium* protocorms were members of *Tulasnella*. Additionally, Zi et al. (2014) reported that *Epulorhiza*, which is an anamorphic state in the genus *Tulasnella*, was found in the protocorms of *Dendrobium*. Furthermore, Athipunyakom et al. (2004) showed that *Trichosporiella multisporum* was found in the roots of *Paphiopedilum*.




Fig. 4. Morphology of non-mycorrhizal fungi: (A) *Fusarium* sp. colony on PDA; (B) Slightly curved macroconidia of *Fusarium* sp. (white arrow); (C) *Beauveria* sp. on PDA; (D) Globose to ellipsoid conidiospore of *Beauveria* sp. (white arrow).

Group II consisted of fungal isolates identified as *Beauveria* sp. and *Fusarium* sp. These fungal isolates were reported from orchid protocorms (Matsubara et al., 2012). Although they were found inside orchid roots, they may be casual, non-mycorrhizal endophytes. Some of these fungi have been previously reported in numerous white rot fungi and are capable of phytopathogenicity (Porras-Alfaro and Bayman, 2011; Quesada-Morage et al., 2014). In another study, a strain of *Fusarium* with orchid seeds yielded a positive result when tested with triphenyl tetrazolium chloride (TTC) and acid fuchsin (AC), suggesting its role in stimulating germination (Vujanovic et al., 2000). However, it is difficult to designate exact ecological roles to these isolates, because the orchids may have different relationships—ranging from mutualistic to slightly pathogenic—at different life stages. More detailed physiological studies will be needed to describe their ecological roles.

These studies provide a better understanding of proper orchid habitats for successful transplanting to the field. More information on orchid fungi ecology is required to accomplish orchid conservation. Therefore, a screening test for fungal compatibility is crucial for symbiotic seed germination. The *in situ* seed baiting technique is a powerful tool for both the production of mycobiont-infected seedlings for use in plant reintroduction and for fungal specificity within the Orchidaceae. Further research should focus on

improving the efficiency of *D. friedericksianum* *in situ* seedling development such as by mass propagation, transplant success to the nursery and then to natural habitats.

This study presents invaluable information on orchid ecology and the endophytic fungi made up of orchid mycorrhizal fungi and non-mycorrhizal fungi associated with *D. friedericksianum* in the natural habitat. The *in situ* seed baiting method was found to be an effective means of detection of compatible mycorrhizal fungi for enhancing orchid seed germination and species-specificity.

Conflict of Interest

No conflicts of interest influenced this research.

Acknowledgments

This work was funded with a grant from the Agricultural Research Development Agency (Public Organization), Bangkok, Thailand, CRP5205010090. The authors would like to express their gratitude to all those who assisted in the field. An anonymous reviewer supplied valuable suggestions for improvement of the text, for which we are thankful.

References

- Arditti, J., Ghani, A.K.A. 2000. Numerical and physical properties of orchid seeds and their biological implications. *New Phytol.* 145: 367–412.
- Athipunyakom, P., Manoch, L., Piluek, C. 2004. Isolation and identification of mycorrhizal fungi from eleven terrestrial orchids. *Kasetsart J. Nat. Sci.* 38: 216–228.
- Bandoni, R.J. 1979. Safranin O as a rapid nuclear stain for fungi. *Mycologia* 71: 873–874.
- Bidartondo, M.I., Read, D.J. 2008. Fungal specificity bottlenecks during orchid germination and development. *Mol. Ecol.* 17: 3707–3716.
- Brundrett, M.C., Scade, A., Batty, A.L., Dixon, K.W., Sivasithamparam, K. 2003. Development of *in situ* and *ex situ* seed baiting techniques to detect mycorrhizal fungi from terrestrial orchid habitats. *Mycol. Res.* 107: 1210–1220.
- Chen, J., Wang, H., Guo, S.X. 2012. Isolation and identification of endophytic and mycorrhizal fungi from seeds and roots of *Dendrobium* (Orchidaceae). *Mycorrhiza* 22: 297–307.
- Chutima, R., Dell, B., Lumyong, S. 2011. Effects of mycorrhizal fungi on symbiotic seed germination of *Pecteilis susanna* (L.) Rafin (Orchidaceae), a terrestrial orchid in Thailand. *Symbiosis* 53: 149–156.
- Convention on International Trade in Endangered Species of Wild Fauna and Flora 2014. Checklist of CITES Species. Available from: <http://www.checklist.cites.org/#/en> [Sourced: 19 May 2012].
- Currah, R.S., Zettler, I.W., McInnis, T.M. 1997. *Epulorhiza inquilina* sp. nov. from *Plantanthera* (Orchidaceae) and a key to *Epulorhiza* species. *Mycotaxon* 31: 335–342.
- Hossain, M.M. 2011. Therapeutic orchids: traditional uses and recent advances-an overview. *Fitoterapia* 82: 102–140.
- Leitch, I.J., Kahandawala, I., Suda, J., Hanson, L., Ingrouille, M.J., Chase, M.W., Fay, M.F. 2009. Genome size diversity in orchids: consequences and evolution. *Annu. Bot.* 104: 469–481.
- Matsubara, T., Yoneda, M., Ishii, I. 2012. Fungal isolate “KMI” is a new type of orchid mycorrhizal fungus. *Am. J. Plant Sci.* 3: 1121–1126.
- McKendrick, S.L., Leake, J.R., Taylor, D.L., Read, D.J. 2000. Symbiotic germination and development of myco-heterotrophic plants in nature; ontogeny of *Corallorhiza trifida* and characterization of its mycorrhizal fungi. *New Phytol.* 145: 523–537.
- Nontachaiyapoom, S., Sasirat, S., Manoch, L. 2010. Isolation and identification of *Rhizoctonia*-like fungi from roots of three orchid genera, *Paphiopedilum*, *Dendrobium*, and *Cymbidium*, collected in Chiang Rai and Chiang Mai provinces of Thailand. *Mycorrhiza* 20: 459–471.
- Nontachaiyapoom, S., Sasirat, S., Manoch, L. 2011. Symbiotic seed germination of *Grammatophyllum speciosum* Blume and *Dendrobium draconis* Rchb.f., native orchid of Thailand. *Sci. Hort.* 130: 303–308.
- Porrás-Alfaro, A., Bayman, P. 2011. Hidden fungi, emergent properties: endophytes and microbiomes. *Annu. Rev. Phytopathol.* 49: 291–315.
- Prasertsivivatna, S., Koolpluksee, M. 2011. The study on optimization of growth conditions for *Dendrobium frederickianum* Rchb.f. seedlings in aseptic culture. *J. Agric. Technol.* 7: 739–749.
- Quesada-Morage, E., Lopez-Diaz, C., Landa, B.B. 2014. The hidden habit of the entomopathogenic fungus *Beauveria bassiana*: first demonstration of vertical plant transmission. *PLoS ONE* 9 (2): e89278. <http://dx.doi.org/10.1371/journal.pone.0089278>.
- Rasmussen, H.N., Rasmussen, F.N. 2009. Orchid mycorrhiza: implications of a mycophagous life style. *Oikos* 118: 334–345.
- Rasmussen, H.N., Whigham, D.F. 1993. Seed ecology of dust seeds *in situ*: a new study technique and its application in terrestrial orchids. *Am. J. Bot.* 80: 1374–1378.
- Roberts, P. 1999. *Rhizoctonia*-forming Fungi: a Taxonomic Guide. Royal Botanic Gardens, Kew, UK.
- Sneh, B., Burpee, L., Ogoshi, A. 1991. Identification of *Rhizoctonia* Species. American Phytopathological Society Press, St Paul, MN, USA.
- Stewart, S.L., Kane, M.E. 2007. Symbiotic seed germination and evidence for *in vitro* mycobiont specificity in *Spiranthes brevilabris* (Orchidaceae) and its implications for species-level conservation. *In Vitro Cell Dev. Bio. Plant* 43: 178–186.
- Vujanovic, V., St-Arnaud, M., Barabe, D., Thibeault, G. 2000. Viability testing of orchid seed and the promotion of colouration and germination. *Ann. Bot.* 86: 79–86.
- Wang, H., Fang, H.Y., Wang, Y.Q., Duan, L.S., Guo, S.X. 2011. *In situ* seed baiting techniques in *Dendrobium officinale* Kimuraet Migo and *Dendrobium nobile* Lindl.: the endangered Chinese endemic *Dendrobium* (Orchidaceae). *World J. Microbiol. Biotechnol.* 27: 2051–2059.
- Yamato, M., Yagame, T., Suzuki, A., Iwase, K. 2005. Isolation and identification of mycorrhizal fungi associating with an achlorophyllous plant, *Epipogium rosesum* (Orchidaceae). *Mycoscience* 46: 73–77.
- Zi, X.M., Sheng, C.L., Goodale, U.M., Gao, J.Y. 2014. *In situ* seed baiting to isolate germination-enhancing fungi for an epiphytic orchid. *Dendrobium aphyllum* Orchid. *Mycorrhiza*. <http://dx.doi.org/10.1007/s00572-2014-0565-8>.