



Research article

Ant community composition in urban areas of Bangkok, Thailand

Sasitorn Hasin^a, Wattanachai Tasen^{b,c,*}^a *Innovation of Environmental Management, College of Innovative Management, Walaya Alongkorn Rajabhat University Under the Royal Patronage, Pathum Thani 13180, Thailand*^b *Center for Advanced Studies in Tropical Natural Resources, NRU-KU, Kasetsart University, Bangkok 10900, Thailand*^c *Department of Forest Biology, Faculty of Forestry, Kasetsart University, Bangkok 10900, Thailand*

Article Info

Article history:

Received 13 June 2019

Revised 27 November 2019

Accepted 23 December 2019

Available online 30 October 2020

Keywords:

Ant communities,

Nesting site,

Pest ant,

Urban ant species,

Urban areas

Abstract

The impact of urbanization on biodiversity is well-documented and ant community composition can be a bio-indicator in habitat disturbance. This necessitates research to understand the diversity and abundance of ants in urban settings. Consequently, ant diversity and abundance were examined using a food baits method and direct sampling and the results were compared among three different habitat types in urban areas in Bangkok city, Thailand. Of the 67 ant species identified from six sub-families in the urban area, five were dominant ant species that had high levels of frequency of occurrence (FO; > 70%). The results indicated that ant richness decreased from green areas to commercial areas. The greatest abundance of ants was in commercial areas where high FO values were recorded for *Trichomyrmex destructor* (100%), *Paratrechina longicornis* (93%), *Tapinoma melanocephalum* (92%), *Monomorium pharaonis* (79%) and *Solenopsis geminata* (71%). The findings revealed that a difference in habitat type in the city had a negative impact on ant diversity and abundance, and a difference in nesting habitat for native ant species was identified for each habitat type. While urbanization might have a positive impact on the abundance of invasive ant species, it was concluded that six species of urban ants (*T. destructor*, *M. pharaonis*, *P. longicornis*, *T. melanocephalum*, *S. geminata* and *Tetraponora rufonigra*) might have become abundant pests in urban areas in Thailand. More research is required to examine the impact of each dominant ant species and their nesting habits in relation to different stages of land development.

Introduction

Land transformations during the process of urbanization affect the habitat structures and resource of native biodiversity (Czech, 2004). Land development such as houses, buildings and yards may occur as a result of increasing human population and human construction. This change results in the loss of local habitats, habitat degradation and increased habitat fragmentation, which have well-documented effects

on biodiversity (McKinney, 2006; Clarke et al., 2008). Numerous studies have reported the decline of native biodiversity due to urbanization (Heterick et al., 2000; Sanford et al., 2008; Vepsäläinen et al., 2008) and the eventual extinction of certain species (Lessard and Buddle, 2005). For example, urbanization has negative effects on species richness and community composition, including those of native plants (Walker et al., 2009), birds (Aronson et al., 2014), amphibians (Riley et al., 2005; Scheffers and Paszkowski, 2012), bats (Coleman and Barclay, 2012) and arthropods (Yamaguchi, 2004; Hartley et al., 2007). Several studies have suggested that urbanization

* Corresponding author.

E-mail address: fforwet@ku.ac.th (W. Tasen)

online 2452-316X print 2468-1458/Copyright © 2020. This is an open access article, production and hosting by Kasetsart University of Research and Development institute on behalf of Kasetsart University.

<https://doi.org/10.34044/j.anres.2020.54.5.07>

and human activities can also affect the activity, reproduction and survival of native animals (George and Crooks, 2006; Villaseñor et al., 2014). On the other hand, urbanization may increase the diversity of alien and invasive species of plants (Nock et al., 2013) and insects (Yamaguchi, 2004; Kamura et al., 2007; Mauro et al., 2007; Catterall, 2009; Menke et al., 2010). Urbanization has spread rapidly in present times, necessitating planning efforts in urban development to conserve urban biodiversity. Thus, it is important to understand how habitat development affects biodiversity.

Ants are one of the excellent bioindicators for studying land-use planning and conservation efforts (Andersen, 2000; Underwood and Fisher, 2006). Since ants are the most abundant and diverse group easily found in any type of habitat (Hölldobler and Wilson, 1990; Bolton, 1994), they can respond quickly to environmental change or human-altered habitats (Stringer et al., 2009). In addition, ants are sensitive to habitat disturbance conditions at the community level (Andersen, 2000) and the negative impacts of ants are well documented, especially invasive ants (Lach and Hooper-Bui, 2010; Rowles and Silverman, 2010). Many studies have reported habitat destruction and its effect on ant community composition (McKinney, 2008). For example, the size, age and shape of a park affected ant diversity and population (Yamaguchi, 2004; Carpintero and Reyes-López, 2013). The urban edge is associated with changes in ant species richness and community composition (Lessard and Buddle, 2005; Pacheco and Vasconcelos, 2007). Uno et al., (2010) reported that vegetation factors (patch size, number and size of trees, leaf litter, amount of concrete and buildings) correlated with differences in ant species composition. For these reasons, ants are useful organisms for examining the effect of habitat development on the biodiversity of urban habitats. The aims of the current study were to examine ant diversity and abundance in urban areas, and to determine the dominant ant species in such an urban ecosystem. The ant species likely to become pests to humans were also discussed.

Materials and Methods

Study site

Fourteen study sites were located in the Bangkok Metropolitan Region (BMR), which is considered a center of human activity (Table 1) as Bangkok is the capital city of Thailand. It is located between 13°45'N and 100°31'E, 2.31 m above sea level and its total area is 1,568.7 km². The current population is over 8 million and the average population density is 3,634 person/km² (Bangkok Geographic Information Technology Center, 2018). Bangkok is the political, social and economic center of Thailand and is surrounded by industrial, commercial and residential areas. In general, the climate in the BMR is influenced by the northeast and southwest monsoons. The southwest monsoon brings heavy rainfall from mid-May to mid-October while the northeast monsoon brings a cool and dry climate from mid-October to February (Limjirakan et al., 2012).

This study was conducted from 2010 to 2012. Sample collection was done separately during the dry season (between November and April) and the wet season (May and October). The study was conducted in three different habitat areas (commercial, residential and green areas) within the 14 study sites in Bangkok and the surrounding provinces. Each study site was classified into three different urban area types based on human activity (Table 1). Commercial area (CM) was defined as an area with buildings for industrial and commercial activities, residential area (RD) as where a group of people live in the same area and green area (GA) as a fragmented green space including gardens, yards, and city parks that is surrounded by commercial and residential areas.

Table 1 Sample site location details

District	Sample site	Coordinates	Urban area type
Bang Khen	Kasetsart University	13°50'51.89"N, 100°34'15.02"E	CM, RD, GA
Chatuchak	Mho Chi bus terminal	13°48'9.30"N; 100°33'13.80"E	CM, RD, GA
	Jatujak Matket,	13°49'43"N, 100°33'35"E	CM, RD, GA
	Queen Sirikit Public Park,	13°48'24" N, 100°33'0"E	CM, RD, GA
	Vachirabenjatat Public Park	13°48'29"N, 100°33'20"E	CM, RD, GA
Klong Toei	Benchasiri Public Park	13° 43'49.22"N, 100°34'2.35"E	CM, RD, GA
Lat Krabang	Suvarnabhumi Airport	13°41'33"N, 100°45'0"E	CM, RD, GA
	Hua Takhe Railway Station	13°43'41.16"N, 100°46'55.92"E	CM, RD, GA
Phatum Wan	Lumpini Public Park	13° 44'36.79"N, 100°32'33.56"E	CM, RD, GA
Phra Nakhon	Sanam Luang	13°45'18"N, 100°29'35"E	CM, RD, GA
	Saran Rom Public Park	13°44'53.71"N, 100 29'42.8"E	CM, RD, GA
Pom Prap Sattru Phai	Shopping mall for import and expofirt of clothes	13.758°N, 100.513°E	CM, RD, GA
Rat burana	Chalermprakiat Public Park	13°40'56"N 100°30'20"E	CM, RD, GA
Ratchathewi	San Ti Pap Public Park	13°45'32"N 100°32'04"E	CM, RD, GA

CM = commercial area; RD = residential area; GA = green area.

Ant sampling design

In each study area, three line transects each 50 m long and 20 m apart were set up for the ant survey. The ants were collected using a food baits method. Baits were made of 5 cm × 5 cm white laminated cards with approximately 2 g of mixed food placed in the center. The food was a mixture of honey, tuna and crushed beans. In total, 10 baits were sampled per transect 5 m away from each other and were left undisturbed for 30 min. The outdoor baits were placed far from trees, garbage and pedestrian walkways and far from garbage for indoor sampling. Any ant that became attached to the bait was collected for identification in the laboratory. The number of individuals was counted for each species.

Direct sampling was used for extensive observation of ant diversity and the location of their nesting sites (tree stratum, above ground, below ground) for each sampling point (Bestelmeyer and Casanova, 2010). At each site, 20 sampling points were systematically established 10 m apart from each other. Ant species and their nesting sites were observed over a 10 min period within 2 m around each point. Nesting sites were identified by the extremely high number of ant workers swarming at their nest entrance hole or cavity or the presence of a queen and brood. All entrance holes or cavities of each species were counted, so that nest entrance numbers could be used to define suitable nesting sites for each ant species. The nesting sites were classified into five types: nesting on trees and lower vegetation (NT), inside or under the human constructions (UW), under or in the leaf litter layer (UT), nesting inside decaying logs (UL) and nesting in the ground (UG).

Ant identification

All ant samples were stored in 95% ethyl alcohol for identification using the available taxonomic keys of Bolton (1994), Hölldobler and Wilson (1990) and Bolton et al. (2006). Later, ants were identified to species or morphospecies level by reference to the ant collection in the Insect Collection at the Department of National Parks, Wildlife and Plant Conservation (DNP), Bangkok, Thailand and a reliable digital resource (<http://www.antweb.org> and <http://www.antbase.de>). Questionable specimens were sent to an expert for confirmation to the species level. Vouchers were deposited in the Insect Collection at the DNP. The general character of ants was characterized based on functional groups of ant to at least the generic level based on Andersen (2000), and invasive ant species were classified according to the literature by McGlynn (1999), Holway et al., (2002) and Office of Natural Resources Environmental Policy and Planning (2009).

Data analyses

Species richness was determined by the total number of ant species pooled from the food baits method and direct sampling at each site. The frequency of occurrence of each species was analyzed for each habitat site using the presence or absence of the ant species in each habitat type. The frequency of occurrence of each species was

calculated separately for each habitat type. Ants were divided into three groups using the frequency of occurrence values (Bourmas, 2005; Hasin, 2008), where a value in the range 0–35% was considered as an uncommon ant species, a value in the range 36–70% was considered as a common ant species and a value in the range 71–100% was considered as the dominant ant species. Abundance was defined as the number of individual workers visiting bait stations.

A univariate analysis of variance (ANOVA) was used to test the effects of habitat type and season on species richness and abundance, with habitat area and season as explanatory variables. All pairwise comparisons were made based on Bonferroni post-hoc tests where the differences were considered significant at $p < 0.05$. Normality and homoscedasticity of the data were confirmed prior to the analyses using the Shapiro-Wilk and Levene tests, respectively. All data were log-transformed to reduce heteroscedasticity prior to analysis. All univariate statistical analyses were performed with PASW ver. 20.0.0 for Windows (SPSS Inc., Chicago, IL, USA).

Results

Community composition of ants

In total, 67 species representing six sub-families and 39 genera were collected (Table 2). The highest richness (67 species) was observed in the green areas, followed by residential areas (29 species) and commercial areas (17 species). A comparison of species richness data from both the food baits method and direct sampling showed significant differences among habitat areas ($F_{2,78} = 69.90$; $p < 0.001$). The average ant species richness was higher in the green areas (28.7 ± 2.6 SE) than in the commercial (5.2 ± 0.3 SE) and residential areas (6.7 ± 0.5 SE) (post-hoc: $p < 0.001$; Fig. 1A). However, ant species richness was not affected significantly by dry and wet seasons ($F_{1,78} = 0.17$; $p = 0.69$), and there was no interaction between study area and season ($F_{2,78} = 0.31$; $p = 0.73$).

A comparison of the ant abundance data from the food baits methods showed significant differences among habitat areas ($F_{2,78} = 5.39$; $p < 0.05$), where the average ant abundance was significantly higher in commercial areas (451.8 ± 32.3 SE) than residential (370.6 ± 36.8 SE) and green areas (311.1 ± 24.4 SE; post-hoc: $p < 0.001$; Fig. 1B). There was no significant difference between the dry and wet seasons ($F_{1,78} = 1.09$; $p = 0.30$), and no interaction between study area and season ($F_{2,78} = 0.20$; $p = 0.82$).

Dominant urban ants

Five ant species dominated the urban areas, namely, *Trichomyrmex destructor* (100%), *Paratrechina longicornis* (93%), *Tapinoma melanocephalum* (92%), *Monomorium pharaonis* (79%) and *Solenopsis geminata* (71%), as shown in Table 2. There were slight differences in frequencies of occurrence of some ant species within the study areas, with *P. longicornis* having higher frequency values in the green areas, and lower in the commercial and residential areas (Table 2).

Table 2 List of ant species sampled and percentage frequency of occurrence in three habitat types and within all habitats combined

Subfamily / species ¹⁾	Direct sampling			Food Bait			F _{habitat} (n=28)			F _{total} (n=84)	Nest site
	CM	RD	GA	CM	RD	GA	CM	RD	GA		
Cerapachyinae											
<i>Cerapachys longitarsus</i> Mayr	-	-	+	0	0	0	0	0	7	3	not found
Dolichoderinae											
<i>Iridomyrmex anceps</i> (Roger)	-	+	+	0	0	102	0	4	36	14	UG
<i>Ochetellus glaber</i> (Mayr)	-	-	+	0	0	0	0	0	18	6	not found
<i>Tapinoma indicum</i> Forel	-	-	+	0	0	3	0	0	18	6	not found
<i>Tapinoma melanocephalum</i> (Fabricius) ¹	+	+	+	2,451	1,582	1,517	86	89	100	92	UG, UW
<i>Technomyrmex butteli</i> Forel	-	-	+	0	0	0	0	0	36	12	UL, UT
<i>Technomyrmex albipes</i> Smith ¹	-	-	+	0	0	98	0	0	11	4	UL, UT
<i>Technomyrmex kraepelini</i> Forel	-	-	+	0	0	24	0	0	14	5	UL, UT
Formicinae											
<i>Anoplolepis gracilipes</i> ¹	-	+	+	0	0	124	0	7	61	23	UL, UT
<i>Camponotus rufoglaucus</i> (Jerdon)	-	+	+	0	2	71	0	4	54	19	UG
<i>Camponotus</i> sp.1	+	+	+	0	0	0	29	50	61	47	UW, UL, UT
<i>Camponotus</i> sp.2	-	-	+	0	0	0	0	0	11	5	not found
<i>Camponotus</i> sp.3	-	-	-	0	0	13	0	0	14	5	not found
<i>Oecophylla smaragdina</i> Fabricius	+	+	+	0	65	44	7	32	68	36	NT
<i>Paratrechina longicornis</i> Latreille ¹	+	+	+	2,132	1,361	807	89	89	100	93	UW,UL,UT,UG
<i>Nylanderia</i> sp.1	-	+	+	0	258	89	0	7	7	3	UW, UL, UT
<i>Nylanderia</i> sp.2	-	-	+	0	0	8	0	0	50	19	not found
<i>Nylanderia</i> sp.3	-	+	+	0	10	13	0	7	21	8	not found
<i>Nylanderia</i> sp.4	-	-	+	0	0	0	0	0	64	22	not found
<i>Plagilepis</i> sp.1	-	-	+	0	0	17	0	0	29	10	not found
<i>Plagilepis</i> sp.2	-	-	+	0	0	33	0	0	21	8	not found
<i>Polyrhachis dives</i> Smith	-	+	+	0	12	5	0	0	36	12	not found
<i>Polyrhachis laevissima</i> Smith	-	-	+	0	0	0	0	0	79	27	not found
<i>Polyrhachis proxima</i> Roger	-	-	+	0	0	3	0	0	43	15	not found
Myrmicinae											
<i>Cardiocondyla emeryi</i> Forel	+	+	+	55	21	18	4	11	96	37	not found
<i>Cardiocondyla nuda</i> (Mayr)	+	+	+	0	6	16	4	7	96	36	not found
<i>Cardiocondyla wroughtonii</i> (Forel)	-	-	+	0	0	0	0	0	4	2	not found
<i>Cataulacus granulatus</i> Latreille	-	-	+	0	0	0	0	0	46	16	not found
<i>Crematogaster rogenhoferi</i> Mayr	-	-	+	0	0	201	0	0	64	22	NT
<i>Crematogaster</i> sp.1	-	-	+	0	0	42	0	0	25	9	not found
<i>Crematogaster</i> sp.2	-	-	+	0	0	55	0	0	43	15	not found
<i>Crematogaster</i> sp.3	-	-	+	0	0	80	0	0	32	11	not found
<i>Meranoplus bicolor</i> Guérin-Méneville	-	-	+	0	0	95	0	0	54	18	UG
<i>Monomorium chinense</i> Santschi	+	+	+	84	4	18	7	11	36	18	UW,UG
<i>Monomorium floricola</i> (Jerdon)	+	+	+	289	38	119	21	32	86	47	UW,UG,UL
<i>Monomorium pharaonis</i> (Linnaeus) ¹	+	+	+	1,029	308	228	64	75	96	79	UW,UG,UL
<i>Monomorium sechellense</i> Emery	-	-	+	0	0	0	0	0	18	7	not found
<i>Monomorium</i> sp.1	-	-	+	0	0	198	0	0	50	17	not found
<i>Myrmicina</i> sp.1	-	-	+	0	0	0	0	0	14	5	not found
<i>Myrmicina</i> sp.2	-	-	+	0	0	0	0	0	7	3	not found
<i>Paratopula macta</i> Bolton	-	-	+	0	0	0	0	0	7	3	not found
<i>Pheidole bugi</i> Wheeler	+	+	+	43	132	55	21	46	93	54	UW,UL, UG
<i>Pheidole fervens</i> (Smith F.)	+	+	+	11	0	0	4	18	29	17	not found
<i>Pheidole plagiaria</i> (Smith F.)	-	+	+	34	0	39	7	4	71	28	UW, UL, UG
<i>Pheidole</i> sp.1	-	-	+	0	0	0	0	0	14	5	not found
<i>Pheidologeton diversus</i> Jerdon	+	+	+	136	62	495	7	21	57	27	UW,UL, UG
<i>Solenopsis geminata</i> Fabricius ¹	+	+	+	1,680	1,024	461	54	57	100	71	UW, UG
<i>Strumigenys</i> sp.1	-	-	+	0	0	0	0	0	11	4	not found
<i>Strumigenys</i> sp.2	-	-	+	0	0	0	0	0	7	3	not found
<i>Tetramorium bicarinatum</i> (Nylander)	+	+	+	0	0	0	7	14	50	24	not found
<i>Tetramorium lanuginosum</i> Mayr	+	+	+	102	62	105	4	36	100	47	UG
<i>Tetramorium simillimum</i> Smith	-	-	+	0	46	56	0	14	43	19	UG

Table 2 Continued

Subfamily / species ¹⁾	Direct sampling			Food Bait			F _{habitat} (n=28)			F _{total} (n=84)	Nest site
	CM	RD	GA	CM	RD	GA	CM	RD	GA		
<i>Tetramorium smithi</i> Mayr	-	-	+	0	103	12	0	18	64	28	UG
<i>Tetramorium walshi</i> Forel	-	-	+	0	95	12	0	14	29	15	UG
<i>Trichomyrmex destructor</i> (Jerdon) ¹	+	+	+	3,603	3,357	2,003	100	100	100	100	UW,UG,UL
Ponerinae											
<i>Anochetus graeffei</i> Mayr	-	-	+	0	0	0	0	0	32	11	UL, UG
<i>Diacamma rugosum</i> LeGuillou	-	-	+	0	0	17	0	0	50	17	UL, UG
<i>Diacamma vagans</i> Smith	-	-	+	0	3	15	0	4	46	16	UL, UG
<i>Hypoponera</i> sp.1	-	-	+	0	0	0	0	0	32	11	not found
<i>Odontomachus simillimus</i> Smith	-	-	+	0	12	24	0	4	54	18	not found
<i>Pachycondyla astuta</i> Smith	-	-	+	0	0	0	0	0	7	3	not found
<i>Pachycondyla leeuwenhoekii</i> Forel	-	-	+	0	0	0	0	0	14	5	not found
<i>Pachycondyla</i> sp.1	-	-	+	0	0	0	0	0	14	5	not found
<i>Pachycondyla</i> sp.2	-	-	+	0	0	0	0	0	11	4	not found
<i>Pachycondyla</i> sp.3	-	-	+	0	0	0	0	0	14	5	not found
Pseudomyrmecinae											
<i>Tetraponera allaborans</i> (Walker)	-	-	+	0	0	0	0	0	11	4	not found
<i>Tetraponera rufonigra</i> Jerdon	-	-	-	0	17	67	0	21	93	39	NT
Total species richness	16	23	65	14	24	42					
Total abundance				11,403	8,826	7,402					

F_{habitat} = three different habitats: CM = commercial area; RD = residential area; GA = green area; F_{total} = within all habitats combined; five nest site abbreviations: NT = on trees and lower vegetation; UW = inside or under human constructions; UT = nesting inside or under dead and living plant material and around base of trees; UL = nesting under or inside leaf litter and decaying logs; UG = nesting in soil.

¹ Invasive pest species were documented using Invasive Species Specialist Group (2010).




Fig. 1 Average values (\pm SE): (A) species richness; (B) abundance of ants, where CM = commercial area; RD = residential area; GA = green area, different letters represent significant differences among habitat sites (post-hoc test; $p < 0.001$) and bars with the same lowercase letter are not significantly different.

The dominant ant species differed within habitat areas as indicated by their frequency of occurrence values ($FO_{habitat}$) of at least 71% (Table 2). Dominant species in the commercial areas were *T. destructor*, *P. longicornis* and *T. melanocephalum*, and in the residential areas were *M. pharaonis*, *T. destructor*, *P. longicornis* and *T. melanocephalum*. Higher richness of the dominant ant species was recorded in the green areas with 11 species. Five of these ant species, namely, *T. destructor*, *P. longicornis*, *T. melanocephalum*, *S. geminata* and *T. lanuginosum*, had large values of frequency of occurrence of 100%. A comparison of the abundance of these ant species showed that ant abundance was significantly different among habitats (one-way ANOVA; $p < 0.01$;

Fig. 2). The mean abundance of *T. melanocephalum* and *P. longicornis* was significantly higher in both the commercial and residential areas than in the green areas (post-hoc: $p < 0.001$). The species *M. pharaonis* and *S. geminata* both had higher abundance values in the residential areas than in the commercial and green areas (post-hoc: $p < 0.05$). There were no differences among habitat types for *T. destructor* ($p > 0.05$).

Interestingly, in the urban areas, invasive ant species were represented by five species of dominant ant, namely *T. destructor*, *P. longicornis*, *T. melanocephalum*, *M. pharaonis* and *S. geminata*, having greater than 50% frequency of occurrence values within all habitats combined (FO_{total} ; Table 2).

Nesting site preference

In total, 29 ant species were found at five sites. The ants had a considerable range in nesting site preferences. Most of the ant species constructed their nests outdoors, particularly in the ground (UG) with 19 species (43% of the total in this observation), inside or under dead or living plant material and around tree bases (NT) with 10 species (24%), inside or under human constructions (UW) with 9 species (21%), under or inside leaf litter and decaying logs (UL) with 3 species (6.9%); and on trees and lower vegetation (UT) with 2 species (5%), as shown in Table 2. Of the studies of the five nest sites, eight ant species were present at only one nesting site and most of the ant species were present at 2–4 nesting sites, excluding the trees (Table 2).




Fig. 2 Average values (\pm SD) abundance of common pest ant species for each study areas from urban ecosystems, where CM = commercial area; RD = residential area; GA = green area and different lowercase letters indicate highly significant differences ($p < 0.01$) within species.

The results clearly showed that the ant diversity and abundance in urban areas were affected by habitat type ($p < 0.001$), and not by season ($p > 0.05$). A larger number of ant species was found in green areas (67 species) compared to the other areas, with all identified species being found in green areas. This suggested that human disturbance of habitat had a negative impact on the number of ant species in urban areas. Similar results were reported by Pacheco and Vasconcelos (2007) and Philpott et al. (2010) who indicated that ant species richness in urban ecosystems declined from parks at urban edges to inner city squares and also around human construction. A larger number of species were found in green areas. One reason for this may have been that the variety in habitat, food and nesting sites within green areas provided a more suitable living environment for ants than did the other areas. Ant community composition is more diverse in green areas especially for species of native ants, which are also the most sensitive regarding habitat disturbance (Menke et al., 2010).

A significant difference was found in ant abundance among habitat types. The commercial areas had greater ant abundance compared to the other two habitats (Fig 1B). However, the results clearly showed that the difference in ant abundance in the study areas may have been related to ant species as there was a higher species abundance in the commercial area than for the other areas, especially for the five dominant ant species of *T. destructor*, *P. longicornis*, *T. melanocephalum*, *M. pharaonis* and *S. geminata* (Salyer et al., 2014; Vonshak and Gordon, 2015).

The results showed that the ant species *T. destructor*, *P. longicornis*, *T. melanocephalum*, *M. pharaonis* and *S. geminata* were the dominant ant species in urban areas. These results were similar to those in CABI (2020), which reported that these ant species have spread widely across Southeast Asia, where there has been higher abundance of them recorded in human habitation. Interestingly, these ant species are members of the list of recognized invasive ant species which are able to adapt and live in human structures and buildings and it is probable that a continuum of anthropogenic activity in urban areas by human disturbance has had a positive effect on the abundance of these five invasive ant species (Lessard et al., 2009; Stringer et al., 2009; Buczkowski and Richmond, 2012; Vonshak and Gordon, 2015).

The five dominant urban ant species in the current study have been identified as household pest that negatively impact the economy and human health (Lee and Robinson, 2001; Eow and Lee, 2007; Man and Lee, 2012). They are characterized by a large number of workers when foraging and their abundance is closely associated with human activities (Schultz and McGlynn, 2000). The current results showed that these dominant urban ants could build their nests in all human environments including structures and buildings in urban areas (Passera 1994; Holway et al., 2002). Moreover, it is possible to classify those ant species as urban bio-indicators that can easily be found in high abundance and can adapt their nesting habits to various human environments.

It was concluded that six species of urban ant (*T. destructor*, *M. pharaonis*, *P. longicornis*, *T. melanocephalum*, *S. geminata*, *T. rufonigra*) might have become considerable pests in the human environment in Thailand, not only due to their variety of nesting sites but also because of their high population numbers. This conclusion was similar to those in previous studies. For example, three species (*T. destructor*, *M. pharaonis*, *S. geminata*) were reported to be ant pests in human environments in Asia (in houses and buildings) by gnawing holes in fabrics and rubber goods and damaging polyethylene cables (Harris et al., 2005). In addition, they can cause damage and threaten human health. For example, *P. longicornis*, *M. pharaonis* and *T. melanocephalum* can transport pathogenic microorganisms, resulting in bacterial infection in hospitals (Moreira et al., 2005; Pantoja et al., 2009; Castro et al., 2015). Some species are aggressive and have painful stings for humans and domestic pets bitten in particular by *S. geminata* and *T. rufonigra* (Piromrat et al., 2008; Potiwat and Sitcharungsi, 2015). Normally, a systemic allergic reaction to ant venom and anaphylactic shock result following being stung by *S. geminata* and *T. rufonigra* (Harris et al., 2005; Potiwat and Sitcharungsi, 2015). In Thailand, *T. rufonigra* has been reported to have a sting that can cause a serious allergic reaction and lead to human death (Potiwat and Sitcharungsi, 2015).

The current research has provided basic information about the ant species that are considered ant pests and their habitat preference in urban areas in Thailand. Therefore, future work should concentrate on examining the environmental and economic harm and the impact on human health caused by the dominant urban ant species, and also investigate changes in ant pest abundance and diversity in relation to different stages of land development. This information will offer direction for future research concerning household pest management in Thailand in relation to the impacts of climate change.

Conflict of Interest

The authors declare that there are no conflicts of interest.

Acknowledgements

This study was supported by the Higher Education Research Promotion and National Research University Project of Thailand, Office of the Higher Education Commission.

References

- Andersen, A.N. 2000. A global ecology of rainforest ant: functional group in relation to environmental stress and disturbance. In: Agosti, D., Alonso, L.E., Majer, J.D., Schultz, T.R. (Eds). *Ant: Standard Methods for Measuring and Monitoring Biodiversity*. Smithsonian Institution Press. Washington, DC, USA, pp. 25–34.
- Aronson, M.F.J., La Sorte, F.A., Nilon, C.H., et al. 2014. A global analysis of the impacts of urbanization on bird and plant diversity reveals key anthropogenic drivers. *Proc. R. Soc. B.* 281: 20133330. doi:10.1098/rspb.2013.3330.
- Bestelmeyer, B., Casanova, L.R. 2010. Field techniques for sampling ants. In: Lach, L., Parr, C.L., Abbott, K.L. (Eds). *Ant Ecology*. Oxford University Press. New York, NY, USA, pp. 39–41.
- Bangkok Geographic Information Technology Center. 2018. Statistics of Population in Bangkok 2018. Bangkok, Thailand. http://www.bangkokgis.com/gis_information/population/, 3 August 2018.
- Bolton, B. 1994. Identification guide to the ant genera of the world. Harvard University Press. Cambridge, MA, USA.
- Bolton, B., Alpert, G., Ward, P.S., Naskrecki, P. 2006. Bolton's catalogue of ants of the world: 1758–2005. Harvard University Press. Cambridge, MA, USA.
- Bourmas, C. 2005. Species diversity of ants at Huay Khayeng, Thong Pha Phum District, Kanchanaburi Province. M.Sc. thesis, Faculty of Forestry, Kasetsart University. Bangkok, Thailand.
- Buczowski, G., Richmond, D.S. 2012. The effect of urbanization on ant abundance and diversity: A temporal examination of factors affecting biodiversity. *PLoS ONE* 7: e41729. doi:10.1371/journal.pone.0041729.
- CABI. 2020. Invasive Species Compendium. Wallingford, England. <https://www.cabi.org/ISC/>, 14 November 2020.
- Carpintero, S., ReyeS-López, J.L. 2013. Effect of park age, size, shape and isolation on ant assemblages in two cities of Southern Spain. *Entomol. Sci.* 17: 41–51. doi:10.1111/ens.12027.
- Castro, M.M., Prezotoa, H.H.S., Fernandes, E.F., Buenob, O.C., Prezotoa, F. 2015. The ant fauna of hospitals: Advancements in public health and research priorities in Brazil. *Rev. Bras. Entomol.* 59: 77–83. <http://dx.doi.org/10.1016/j.rbe.2015.02.011>.
- Catterall, C.P. 2009. Responses of faunal assemblages to urbanization: global research paradigms and an avian case study. In: McDonnell, M.J., Hahs, A.K., Breuste, J.H. (Eds). *Ecology of Cities and Towns: A Comparative Approach*. Cambridge University Press. Cambridge MA, USA, pp. 129–155.
- Clarke, K.M., Fisher, B.L., LeBuhn, G. 2008. The influence of urban park characteristics on ant (Hymenoptera, Formicidae) communities. *Urban Ecosyst.* 11: 317–334. doi: 10.1007/s11252-008-0065-8.
- Coleman, J.L., Barclay, R.M.R., 2012. Urbanization and the abundance and diversity of prairie bats. *Urban Ecosyst.* 15: 87–102. doi: 10.1007/s11252-011-0181-8.
- Czech, B. 2004. Urbanization as a threat to biodiversity: Trophic theory, economic geography, and implications for conservation land acquisition. In: David, N. (Eds). *Policies for Managing Urban Growth and Landscape Change: A Key to Conservation in the 21st Century* 265. U.S. Department of Agriculture, Forest Service, North Central Research Station. Madison, WI, USA, pp. 8–13.
- Eow, A.G.H., Lee, C.Y. 2007. Comparative nutritional preferences of tropical pest ants, *Monomorium pharaonis*, *Monomorium floricola*, *Monomorium destructor* (Hymenoptera: Formicidae). *Sociobiology* 49: 165–186.
- George, S.L., Crooks, K.R. 2006. Recreation and large mammal activity in an urban nature reserve. *Biol. Cons.* 133: 107–117.
- Harris, R., Abbott, K., Barton, K., et al. 2005. Invasive Ant Pest Risk Assessment Project for Biosecurity New Zealand. Series of Unpublished Landcare Research Contract Reports to Biosecurity New Zealand. Wellington, NZ. http://www.landcareresearch.co.nz/research/biosecurity/stowaways/Ants/ant_pest_risk.asp.
- Hartley, D.J., Koivula, M.J., Spence, J.R., Pelletier, R., Ball, G.E. 2007. Effects of urbanization on ground assemblages (Coleoptera, Carabidae) of grassland habitats in western Canada. *Ecography* 30: 673–684. doi.org/10.1111/j.2007.0906-7590.05199.x
- Hasin, S. 2008. Diversity and Community Structure of Ants at Sakaerat Environmental Research Station, Nakhon Ratchasima Province. M.Sc. thesis, Faculty of Forestry, Kasetsart University. Bangkok, Thailand.
- Heterick, B.E., Casella, J., Majer, J.D. 2000. Influence of Argentine and coastal brown ant (Hymenoptera: Formicidae) invasions on ant communities in Perth gardens, Western Australia. *Urban Ecosyst.* 4: 277–292.
- Hölldobler, B., Wilson, E.O. 1990. *The Ants*. Harvard University Press. Cambridge, MA, USA.
- Holway, D.A., Lach, L., Suarez, A.V., Tsutsui, N.D., Case, T.J. 2002. The causes and consequences of ant invasions. *Annu. Rev. Ecol. Evol. Syst.* 33: 181–233. doi.org/10.1146/annurev.ecolsys.33.010802.150444
- Invasive Species Specialist Group. 2010. The Global Invasive Species Database. International Union for Conservation of Nature, Italy. <http://www.iucngisd.org/gisd/>, 30 October 2020.
- Kamura, C.M., Morini, M.S.C., Figueiredo, C.J., Bueno, O.C., Campos-Farinha, A.E.C. 2007. Ant communities (Hymenoptera: Formicidae) in an urban ecosystem near the Atlantic Rainforest. *Braz. J. Biol.* 67: 635–641. doi.org/10.1590/S1519-69842007000400007
- Lach, L., Hooper-Bui, L.M. 2010. Consequences of ant invasions. In: Lach, L., Parr, C., Abbott, K. (Eds.). *Ant Ecology*. Oxford University Press. Oxford, UK. pp. 261–286.
- Lee, C.Y., Robinson, W.H. 2001. *Handbook of Malaysian Household and Structural Pests*. Pest Control Association of Malaysia. Kuala Lumpur, Malaysia.
- Lessard, J.P., Buddle, C.M. 2005. The effects of urbanization on ant assemblages (Hymenoptera: Formicidae) associated with the Molson Nature Reserve, Quebec. *Can. Entomol.* 137: 215–225. doi.org/10.4039/n04-055
- Lessard, J.P., Fordyce, J.M., Gotelli, N.J., Sanders, N.J. 2009. Invasive ants alter the phylogenetic structure of native communities. *Ecology* 90: 2664–2669. doi.org/10.1890/09-0503.1
- Limjirakan, S., Limsakul, A. 2012. Observed trends in surface air temperatures and their extremes in Thailand from 1970 to 2009. *J. Meteorol. Soc. JPN.* 90: 647–662. doi: 10.2151/jmsj.2012-505
- Man, L.S., Lee, C.Y. 2012. Structure-invading pest ants in healthcare facilities in Singapore. *Sociobiology* 59: 241–249. doi: 10.13102/sociobiology.v59i1.681
- Mauro, D.D., Dietz, T., Rockwood, L. 2007. Determining the effect of urbanization on generalist butterfly species diversity in butterfly gardens. *Urban Ecosyst.* 10: 427–439. doi: 10.1007/s11252-007-0039-2
- McGlynn, T.P. 1999. The worldwide transfer of ants: geographical distribution and ecological invasions. *J. Biogeogr.* 26: 535–548.
- McKinney, M.L. 2006. Urbanization as a major cause of biotic homogenization. *Biol. Conserv.* 127: 247–260.
- McKinney, M.L. 2008. Effects of urbanization on species richness: a review of plants and animals. *Urban Ecosyst.* 11: 161–176.
- Menke, S.B., Guénard, B., Sexton, J.O., Weiser, M.D., Dunn, R.R., Silverman, J. 2010. Urban areas may serve as habitat and corridors for dry-adapted, heat tolerant species; An example from ants. *Urban Ecosyst.* 14: 135–163. doi: 10.1007/s11252-010-0150-7.
- Moreira, D.D.O., Morais, V., da Motta, O.V., Campos-Farinha, A.E.C., Tonhasca, A. 2005. Ants as carriers of antibiotic-resistant bacteria in hospitals. *Neotrop. Entomol.* 34: 999–1006. doi.org/10.1590/S1519-566X2005000600017

- Nock, C.A., Paquette, A., Follett, M., Nowak, D.J., Messier, C. 2013. Effects of urbanization on tree species functional diversity in eastern North America. *Ecosystems* 16:1487–1497. doi: 10.1007/s10021-013-9697-5
- Pacheco, R., Vasconcelos, H.L. 2007. Invertebrate conservation in urban areas: Ants in the Brazilian Cerrado. *Landsc. Urban Plan.* 81: 193–199.
- Pantoja, L.D.M., Filho, R.E.M., Brito, E.H.S., et al. 2009. Ants (Hymenoptera: Formicidae) as carriers of fungi in hospital environments: An emphasis on the genera *Tapinoma* and *Pheidole*. *Ann. Entomol. Soc. Am.* 46: 895–899. doi.org/10.1603/033.046.0423.
- Passera, L. 1994. Characteristics of tramp species. In: Williams, D.F. (Eds) *Exotic Ants: Biology, Impact, and Control of Introduced Species*. Westview Press. Boulder, Colorado, USA. pp. 23–43.
- Philpott, S.M., Perfecto, I., Armbrrecht, I., Parr, C.L. 2010. Ant diversity and function in disturbed and changing habitats. In: Lach, L., Parr, C.L., Abbott, K.L. (Eds). *Ant Ecology*. Oxford University Press. New York, NY, USA, pp. 137–157.
- Piromrat, K., Chinratapisit, S., Trathong, S. 2008. Anaphylaxis in an emergency department: A 2-year study in a tertiary-care hospital. *Asian Pac. J. Allergy Immunol.* 26: 121–128.
- Potiwat, R., Sitcharungsi, R. 2015. Ant allergens and hypersensitivity reactions in response to ant stings. *Asian Pac. J. Allergy Immunol.* 33: 267–275.
- Riley, S.D., Busteed, G.T., Kats, L.B., et al. 2005. Effects of urbanization on the distribution and abundance of amphibians and invasive species in Southern California streams. *Conserv. Biol.* 19: 1894–1907.
- Rowles, A.D., Silverman, J. 2010. Argentine ant invasion associated with loblolly pines in the Southeastern United States: minimal impacts but seasonally sustained. *Environ. Entomol.* 39: 1141–1150. doi: 10.1603/EN10004.
- Salter, A., Bennett, G.W., Buczkowski, G.A. 2014. Odorous house ants (*Tapinoma sessile*) as back-seat drivers of localized ant decline in urban habitats. *PLoS ONE*. 9: e113878. doi:10.1371/journal.pone.0113878.
- Sanford, M., Manley, P.N., Murphy, D.D. 2008. Effects of urban development on ant communities: Implications for ecosystem services and management. *Conserv. Biol.* 23: 131–141. doi: 10.1111/j.1523-1739.2008.01040.x.
- Scheffers, B.R., Paszkowski, C.A. 2012. The effects of urbanization on North American amphibian species: Identifying new directions for urban conservation. *Urban Ecosyst.* 15: 133–147. doi 10.1007/s11252-011-0199-y.
- Schultz, T.R., McGlynn, T.P. 2000. The interactions of ants with other organisms. In: Agosti, D., Majer, J., Alonso, L., Schultz, T.R. (Eds.) *Ants: Standard Methods for Measuring and Monitoring Biodiversity*. Smithsonian Institution Press. Washington, DC, USA, pp. 35–44.
- Stringer, L., Stephens, A., Suckling, D., Charles, J. 2009. Ant dominance in urban areas. *Urban Ecosyst.* 12: 503–514.
- Tam, K.C., Bonebrake, T.C. 2015. Butterfly diversity, habitat and vegetation usage in Hong Kong urban parks. *Urban Ecosyst.* 19: 721–733. doi: 10.1007/s11252-015-0484-2.
- Underwood, E.C., Fisher, B.L. 2006. The role of ants in conservation monitoring: If, when, and how. *Biol. Conserv.* 132: 166–182.
- Uno, S., Cotton, J., Philpott, S.M. 2010. Diversity, abundance, and species composition of ants in urban green spaces. *Urban Ecosyst.* 13: 425–441. doi: 10.1007/s112 52-010-0136-5.
- Vepsäläinen, K., Ikonen, H., Koivula, M.J. 2008. The structure of ant assemblages in an urban area of Helsinki, southern Finland. *Ann. Zool. Fenn.* 45: 109–127. doi: 10.5735/086.045.0203
- Villaseñor, N.R., Driscoll, D.A., Escobar, M.A.H., Gibbons, P., Lindenmayer, D.B. 2014. Urbanization impacts on mammals across urban-forest edges and a predictive model of edge effects. *PLoS ONE* 9: e97036. doi:10.1371/journal.pone.0097036.
- Vonshak, M., Gordon, D.M. 2015. Intermediate disturbance promotes invasive ant abundance. *Biol. Conserv.* 186: 359–367. doi.org/10.1016/j.biocon.2015.03.024
- Walker, J.S., Grimm, N.B., Briggs, J.M., Gries, C., Dugan, L. 2009. Effects of urbanization on plant species diversity in central Arizona. *Front. Ecol. Environ.* 7: 465–470. doi.org/10.1890/080084
- Yamaguchi, T. 2004. Influence of urbanization on ant distribution in parks of Tokyo and Chiba City, Japan I. Analysis of ant species richness. *Ecol. Res.* 19: 209–216.