

EFFECT OF ESSENTIAL OIL OF *Carum carvi* ON GROWTH INHIBITION OF PATHOGENIC BACTERIA

H. Aroiee¹* and M. Mohsenzadeh²

¹Department of Horticulture, Ferdowsi university of Mashhad, Mashhad-Iran.

²Department of Food Hygiene and Technology, Ferdowsi university of Mashhad, Mashhad-Iran.

ABSTRACT

The antimicrobial activities of the essential oils of Iranian caraway "*Carum carvi*" were investigated against *Escherichia coli* ATCC 25922, *Klebsiella pneumonia* ATCC 13883, *Proteus vulgaris* ATCC 13315, and *Pseudomonas aeruginosa* ATCC 27853. The antimicrobial activities of the essential oils were evaluated by disc diffusion method. The results showed that the essential oil of *Carum carvi* inhibited the growth of *E. coli* and *K. pneumonia*. Furthermore, the study suggests that these essential oils can be used as preservatives in foods.

KEYWORDS: *Carum carvi*, Antibacterial activity, *Escherichia coli*, *Proteus vulgaris*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, Essential oils

1. INTRODUCTION

The growing interest in the substitution of traditional food preservatives, both antimicrobials activities and antioxidants, by natural ones has fostered research on screening of plant materials in order to identify new compounds. It has been known since ancient times that medicinal, aromatic and spice plants and their essential oils (active substances) have varying degrees of anti microbial activity [2][5,6][12,13][19-24][26,27]. The plants contain numerous biologically active compounds, many of which have been shown to have anti microbial properties [8]. The major antimicrobial components of these plants and their essential oils are, for example, eugenol in cloves, allicin in garlic, cinnamic aldehyde and eugenol in cinnamon, carvacrol and thymol in oregano and thyme, and vanillin in vanilla beans [19,25]. Furthermore, essential oils of many aromatic plants have been shown to possess antimicrobial activities [7,10]. Essential oils of basil, bay, clove, thyme and rosemary was reported to have bactericidal activities against *Leisteria monocytogenes* and other pathogens [15,17]. Incidences of foodborne illnesses are still a major problem, even in developed countries. In fact, food poisoning is still a threat for both consumers and the food industry despite the use of preserving processes. Meanwhile, consumers are concerned about the safety of foods containing preservatives. Therefore, there has been a growing interest in new and effective techniques to reduce cases of foodborne illnesses. There is considerable interest in the possible use of these compounds as food preservatives, to delay the onset of food spoilage or to prevent the growth of foodborne pathogens.

The aim of the present study was to evaluate the antibacterial activity of medicinal plant essential oil of *Carum carvi* against common pathogens.

2. MATERIALS AND METHODS

2.1 Extraction of essential oil

The plant investigated in this study is *Carum carvi* (caraway) from umbelliferous family. Caraway is a member of the group of aromatic plants characterized by carminative properties, like anise, cumin, dill

* Corresponding author.

E-mail: aroiee_h@yahoo.com, aroiee@ferdowsi.um.ac.ir

and fennel. It is a biennial, with smooth, furrowed stems growing 1 1/2 to 2 feet high, bearing finely cut leaves, and umbels of white flowers which blossom in June. The fruits (seeds) and the oil extracted from the fruits are as a medicinal properties, flavouring in cookery, confectionery and liqueurs. The seeds contain 4-7 % of essential (volatile) oil. The chief constituent of the oil is a hydrocarbon termed Carvone and an oxygenated oil Carvyl. Both fruit and oil possess aromatic, stimulant and carminative properties [28]. The characteristics of the plant are in Table 1. The air dried and crushed seeds of *Carum carvi* were extracted by hydrodistillation in a Clevenger-type apparatus to obtain essential oil. The plant material was placed in water and heated to boil. The essential oil was condensed, collected in a receiver flask, and separated from water by separator container. Then the essential oil was used in the formula as a main material. The contents of essential oil components of *Carum carvi* are shown in Table 2.

Table 1. Scientific characteristics of the plant that used in this experiments.

Name	Botanical name	Family	Part used for essential oil extraction
Black Caraway	<i>Carum carvi</i>	Apiaceae	Fruits (Seeds)

Table 2. The approximate content of essential oil of *Carum carvi* seeds (GC Varian with capillary column)

	Components	%
1	Alpha Pinen	0.19
2	Beta Pinen	0.7
3	Beta Phelandrene	4.5
4	Para Cimen	3
5	Gama Terpinen	10
6	Cumin Aldehyde	20
7	Cumin Alcohol	50

2.2 Microorganisms

The four bacterial strains, *Escherichia coli* ATCC 25922, *Klebsiella pneumonia* ATCC 13883, *Proteus vulgaris* ATCC 13315, and *Pseudomonas aeruginosa* ATCC 27853 were used as test microorganisms in this experiment. The bacteria were incubated on a nutrient agar slant (Stationary culture) for 48 h at 37 °C followed by inoculation in Mueller Hinton Agar medium. The bacteria were supplied by the Department of Microbiology, Medical Science university., Mashhad, Iran.

2.3 Antibacterial assay

Antimicrobial activity was demonstrated using a modification of the method originally described by Kirby-Bauer which is widely used for the antibacterial susceptibility testing [1]. A loopful of bacteria was taken from the stock culture added in 1 ml of brain heart infusion broth (BHI) and incubated at 35 °C. Final cell concentration was 10⁶-10⁷ CFU/ml. One milliliter of this inoculum was added to each plate containing nutrient agar. All the tests were done by placing sterile paper discs on the Mueller Hinton Agar surface previously inoculated with tested microorganisms with various concentrations. Fifty microliter of essential oil at each concentration were added on a paper disc. Distilled water at a concentration of 50 µl/disc was also added as a negative control. Plates were incubated at 37°C for 24 h to observe formation of inhibition zones around the disc.

3. RESULTS AND DISCUSSION

The results of antimicrobial effect of essential oil against tested bacteria are shown in Table 3. Essential oil of *Carum carvi* were ineffective against, *Proteus vulgaris* and *Pseudomonas aeruginosa*. The results showed that the essential oil of *Carum carvi* was effective in controlling of *Klebsiella pneumonia*. The investigations of Valero and Salmeron [25] showed that the essential oil of cinnamon, oregano and

thyme were most effective on controlling of *B. cereus*[25]. Essawi and Srour [11] reported that *Salvia officinalis*, *Teucrium polium*, *Majorana syriaca*, *Thymus origanium*, *Thymus vulgaris*, *Commiphora opobalsamum*, *Foeniculum vulgare*, and *Rosmarinus officinalis* exhibited an antibacterial effect against some of gram-positive and gram-negative bacteria[11]. It has been indicated that the antibacterial activity of some medicinal and aromatic plants are due to different chemical agents in the extracts, including essential oils (specially thymol), flavonoids and triterpenoids and other compounds of phenolic nature or free hydroxyl group, which are classified as active antimicrobial compounds [11,18]. The growth of *Escherichia coli* in comparison of *Klebsiella pneumoniae* was not inhibited by the essential oil, but *Pseudomonas aeruginosa* did not have any reaction to the essential oil. Camporese *et al.* [4] reported that various concentrations of the plant extracts showed activity to some extent against *Escherichia coli* and *Pseudomonas aeruginosa*, while *Aristolochia trilobata* leaves and bark *Syngonium podophyllum* leaves and bark were also active against *Staphylococcus aureus* [4]. A number recent studys demonstrated an antibacterial activity of rosmarinic acid against *E. coli* and *S. aureus* [14]. Sagdic *et al.*[19] reported that among extracts of seven spices, thyme and oregano showed higher activity than cloves, garlic, and vanilla beans on the growth of *Escherichia coli* 0157:H7 [19].Natural substances from plants (specially, medicinal, aromatic and spice plants) have applications in controlling pathogens in foods [3,9]. Since the medicinal plants studied appear to have a broad anti microbial activity spectrum, they could be useful in antiseptic and disinfectant formulations as well as in chemotherapy [16].

4. CONCLUSION

Furthermore, beside the confirmation of the popular uses of medicinal plants, the obtained results demonstrate that these herbal drugs could represent a new source of antimicrobial agents, less expensive than the imported drugs (4).

REFERENCES

- [1] Barry, A.L., and Thornsberry, C. 1985 Susceptibility Tests. Disc Diffusion Test Procedures. In: Lennette Balows, E.H.; Hauster, W.J., and Shadomy, H.J. (Eds.), *Manual of Clinical Microbiology*. Washington, DC, American Society for Microbiology, pp. 978–987.
- [2] Beuchat, L.R., and Golden, D.A.1989 Antimicrobial Occurring Naturally in Food, *Food Technology*, 43 , 134-142.
- [3] Bowles, B.L., and Juneja, V.K.1998 Inhibition of Food-Borne Bacterial Pathogens by Naturally Occurring Food Additives, *Journal of Food Safety*, 18 : 101-112.
- [4] Camporese, A., Balick, M.J., Arvigo, R., Esposito, R.G., Morsellino, N., De Simone, F., and Tubaro, A. 2003 Screening of Anti-Bacterial Activity of Medicinal Plants from Belize (Central America), *Journal of Ethnopharmacology*, 87: 103–107.
- [5] Chaibi, A., Ababouch, L.H., Bllasri, K., Boucetta, S., and Busta, F.F. 1997 Inhibition of Germination and Vegetative Gowth of *Bacillus cereus* T and *Clostridium botulinum* 62A Spores by Essential Oils, *Food Microbiology*, 14 , 161-174.
- [6] Chang, H.W. 1995 Antimicrobial Effect of Spices and Vegetables, *Food Industries (ROC)*, 27, 53-61
- [7] Cosentino, S., Tuberoso, C.I.G., Pisano, B., Satta, M., Mascia, V., Arzedi, E., and Palmas, F.1999 In Vitro Antimicrobial Activity and Chemical Composition of Sardinian *Thymus* Essential Oils, *Lett. Appl. Microbiol.*, 29, 130– 135.
- [8] Cowan, M.M.1999 Plant Products as Anti Microbial Agents, *Clinical Microbiology Reviews*, 12, 564–582.
- [9] Davidson, P.M. 1997 Chemical Preservatives and Natural Antimicrobial Compounds : In : Doyle, M.P. , Beuchat, L.R., Montville, T.J. (Eds.), *Food Microbiology . Fundamental and Frontiers*.

ASM Publications, Washington, DC, pp.520-556.

[10] Elgayar, M., Draughon, F.A., Golden, D.A., and Mount, J.R. **2001** Antimicrobial Activity of Essential Oils from Plants Against Selected Pathogenic and Saprophytic Microorganisms, *J. Food Prot.*, 64, 1019– 1024.

[11] Essawi, T., and Srour, M. **2000** Screening of Some Palestinian Medicinal Plants for Antibacterial Activity, *Journal of Ethnopharmacology*, 70 : 343–349.

[12] Juven, B.J., Kanner, J., Schved, F., and Weisslowicz, H. **1994** Factors That Interact with the Antibacterial Action of Thyme Essential Oil and Its Active Constituents, *Journal of Applied Bacteriology*, 76 , 626-631.

[13] Lachowicz, K.J., Jones, G.P., Briggs, D.R., Bienvenu, F.E., Wan, J., Wilcock, A., and Coventry, M.J. **1998** The Synergistic Preservative Effects of the Essential Oils of Sweet Basil (*Ocimum basilicum* L.) Against Acid Tolerant Food Microflora, *Letters in Applied Microbiology*, 26 , 209-214.

[14] Li, R., Teng, D., Du, G., and Wang, B. **2000** Isolation of Rosmarinic Acid from Callus of *Perilla frutescens* and Studies of Its Inhibition on the Growth of Bacteria and Fungus, *Weishengwuxue Tongbao*, 27, 324–327.

[15] O'Gara, E., Hill, D.J., and Maslin, D.J. **2000** Activities of Garlic Oil, Garlic Powder, and Their Diallyl Constituents Against *Helicobacter Pylori*, *Appl. Environ. Microbiol.*, 66, 2269– 2273.

[16] Olukoya, D.K., Idika, N., and Odugbemi, T. **1993** Antimicrobial Activity of Some Medicinal Plants from Nigeria. *Journal of Ethnopharmacology*, 39, 69–72.

[17] Smith-Palmer, A., Stewart, J., and Fyfe, L. **1998** Antimicrobial Properties of Plant Essential Oils and Essences Against Important Foodborne Pathogens, *Lett. Appl. Microbiol.*, 26, 118–122.

[18] Rojas, A., Hernandez, L., Pereda-Miranda, R., and Mata, R. **1992** Screening for Antimicrobial Activity of Crude Drug Extracts and Pure Natural Products from Mexican Medicinal Plants, *Journal of Ethnopharmacology*, 35, 275–283.

[19] Sagdic, O., Kuscu, A., Ozcan, M., and Ozcelik, S. **2002** Effects of Turkish Spice Extracts at Various Concentrations on the Growth of *Escherichia coli* 0157:H7, *Food Microbiology* , 19: 473 – 480.

[20] Shelef, L.A. **1983** Antimicrobial Effects of Spices, *Journal of Food Safety*, 6 , 29-44.

[21] Sivropoulo, A., Papanikolaou, E., Nikolaou, C., Kokkini, S., Lanaras, T., and Arsenakis, M. **1996** Antimicrobial and Cytotoxic Activities of Origanum Essential Oils, *Journal of Agricultural and Food Chemistry*, 44 , 1202-1205.

[22] Sokmen, A., Jones, B. M., and Erturk, M. **1999** The In Vitro Antibacterial Activity of Turkish Medicinal Plants, *Journal of Ethnopharmacology*, 67 : 79-86.

[23] Tassou, C.C., Drosinos, E.H., and Nychas, G.J.E. **1995** Effects of Essential Oil From Mint(*Mentha piperita*) on *Salmonella enteritidis* and *Listeria monocytogenes* in Model Food Systems at 4 °C and 10 °C, *Journal of Applied Bacteriology*, 78, 593-600.

[24] Ting, E.W.T., and Deibel, K.E. **1992** Sensitivity of *Listeria monocytogenes* to Spices at Two Temperatures, *Journal of Food Safety*, 12 , 129-137.

[25] Valero, M., and Salmeron, M.C. **2003** Antibacterial Activity of 11 Essential Oils Against *Bacillus Ocereus* in Tyndallized Carrot Broth, *International Journal of Food Microbiology* , 85 : 73 – 81.

[26] Wan,J., Wilcock, A., and Coventry, M. J. **1998** The Effect of Essential Oils of Basil on the Growth of *Aeromonas hydrophila* and *Pseudomonas fluorescens*, *Journal of Applied Microbiology*, 84 , 152-158.

[27] Zaika, L.L. **1988** Spices and Herbs: Their Antimicrobial Activity and Its Determination, *Journal of Food Safety*, 9 , 97-118.

[28] <http://www.Botanica.com>, A modern Herbal, Grieve, M. **2006** Caraway, *Botanica.com*