

Efficacy of Mycotoxin Adsorbents on Aflatoxin B1 Decontamination and *in vitro* Rumen Fermentation

Chitraporn Yeanpet¹, Chaiyapas Thamrongyoswittayakul^{1,2*}, Chalong Wachirapakorn³,
Piyada Songsermsakul⁴, Noppadon Somphon² and Chuntira Wongnen³

¹Research Group of Animal Health Technology, Faculty of Veterinary Medicine, Khon Kaen University,
Khon Kaen, 40002, Thailand

²Division of Livestock Medicine, Faculty of Veterinary Medicine, Khon Kaen University, Khon Kaen, 40002, Thailand

³Department of Animal Science, Faculty of Agriculture, Khon Kaen University, Khon Kaen, 40002 Thailand

⁴Department of Toxicology, Faculty of Pharmaceutical Science, Khon Kaen University, Khon Kaen, 40002, Thailand

Abstract

This study was conducted by using an *in vitro* gas production technique to study the effect of aflatoxin B1 and mycotoxins adsorbents on aflatoxin B1 decontamination and efficiency of rumen fermentation. There were eight treatments, that assigned into completely Randomized Design: T1=the control group, T2= the diet plus 1 µg/mg of aflatoxin B1 (AFB1), T3, T4, T5 = the diets supplement 0.5% of each adsorbent (activated charcoal, montmorillonite, *Saccharomyces cerevisiae*) and T6, T7, T8 = the diets plus 1 µg/mg of AFB1 together with 0.5% of each mycotoxin adsorbent, respectively. The results were shown that the gas production from the insoluble fraction and the potential extent of gas production in the diet plus the AFB1 were lower than the control (P<0.05). Moreover, montmorillonite and *S. cerevisiae* were effectively reduced the negative effect of AFB1 on gas production. In contrast, the diet plus AFB1 and diet plus mycotoxin adsorbents did not interfere the *in vitro* dry matter and organic matter digestibilities, volatile fatty acids and ammonia nitrogen (P>0.05). In conclusion, AFB1 had effected and had decreased cumulative gas production. However, montmorillonite and *S. cerevisiae*, as effective adsorbents, could be reduced the adverse effect of AFB1 in this study.

Keywords: Aflatoxin B1, Mycotoxin adsorbent and Rumen fermentation

* Corresponding author: E-mail: chaiyapas@kku.ac.th

ประสิทธิภาพของสารดูดซับพิษจากเชื้อราต่อการจัดการปนเปื้อนของอะฟลาทอกซิน ปี 1 และ การหมักในกระเพาะรูเมนในหลอดทดลอง

จิตรภรณ์ เยียนเพชร¹, ไชยพัศร์ อารงยศวิทยากุล^{1,2*}, ฉลอง วชิราภรณ์³,
ปิยะดา ส่งเสริมสกุล⁴, นพดล สมผล² และ จันทิรา วงศ์เนตร³

¹ กลุ่มวิจัยเทคโนโลยีสุขภาพสัตว์ คณะสัตวแพทยศาสตร์ มหาวิทยาลัยขอนแก่น อำเภอเมือง จังหวัดขอนแก่น 40002

² กลุ่มอายุรศาสตร์ปศุสัตว์ คณะสัตวแพทยศาสตร์ มหาวิทยาลัยขอนแก่น อำเภอเมือง จังหวัดขอนแก่น 40002

³ ภาควิชาสัตวศาสตร์ คณะเกษตรศาสตร์ มหาวิทยาลัยขอนแก่น อำเภอเมือง จังหวัดขอนแก่น 40002

⁴ ภาควิชาพิษวิทยา คณะเภสัชศาสตร์ มหาวิทยาลัยขอนแก่น อำเภอเมือง จังหวัดขอนแก่น 40002

บทคัดย่อ

การศึกษานี้ได้ดำเนินการโดยใช้เทคนิค *in vitro* gas production technique เพื่อศึกษาผลของอะฟลาทอกซิน ปี 1 และสารดูดซับสารพิษจากเชื้อราชนิดต่างๆต่อการลดการปนเปื้อนของสารพิษจากเชื้อราและประสิทธิภาพของการหมัก โดยวางแผนการทดลองแบบสุ่มสมบูรณ์ แบ่งอาหารทดลองออกเป็น 8 ทริตเมนต์ ประกอบด้วย อาหารกลุ่มควบคุม (T1), อาหารเสริมอะฟลาทอกซิน ปี 1 เข้มข้น 1 ไมโครกรัม/อาหาร 1 มก. (T2), อาหารเสริมสารดูดซับที่ระดับร้อยละ 0.5 (ผงถ่านกัมมันต์ (T3), มอนต์โมริลโลไนต์ (T4), ยีสต์ *Saccharomyces cerevisiae* (T5), และอาหารเสริมอะฟลาทอกซิน ปี 1 1 ไมโครกรัม/1 มก. ร่วมกับสารดูดซับชนิดต่างๆ ที่ระดับร้อยละ 0.5 (T6, T7, T8) ตามลำดับ ผลการศึกษาพบว่า ปริมาณแก๊สที่ผลิตจากส่วนที่ละลายยาก และปริมาณผลผลิตแก๊สที่ผลิตได้ทั้งหมดในอาหารที่มีของอะฟลาทอกซิน ปี 1 ต่ำกว่าในอาหารกลุ่มควบคุม ($P < 0.05$) นอกจากนี้ ยังพบว่ามอนต์โมริลโลไนต์ และยีสต์ *S. cerevisiae* มีประสิทธิภาพในการลดผลกระทบของอะฟลาทอกซิน ปี 1 ต่อการผลิตแก๊ส ในทางตรงข้ามอาหารที่เติมอะฟลาทอกซิน ปี 1 และอาหารเสริมสารดูดซับสารพิษจากเชื้อราไม่มีผลกระทบต่อ การย่อยได้ของวัตถุดิบและการย่อยได้ของอินทรีย์วัตถุในหลอดทดลองปริมาณกรดไขมันที่ระเหยง่าย และแอมโมเนีย - ไนโตรเจน ($P > 0.05$) สรุปได้ว่า การศึกษานี้พบว่าอะฟลาทอกซิน ปี 1 มีผลต่อกระบวนการหมักในหลอดทดลอง และการเสริมมอนต์โมริลโลไนต์ และยีสต์ *S. cerevisiae* ที่ระดับร้อยละ 0.5 เป็นสารดูดซับสารพิษจากเชื้อราพบว่ามีประสิทธิภาพสามารถลดผลไม่พึงประสงค์ของอะฟลาทอกซิน ปี 1 ได้

คำสำคัญ: อะฟลาทอกซิน ปี 1 สารดูดซับสารพิษจากเชื้อรา และ การหมักในกระเพาะรูเมน

* ผู้เขียนให้ติดต่อ: E-mail: chaiyapas@kku.ac.th

Introduction

Aflatoxins (AFs) are secondary metabolites commonly produced by *Aspergillus flavus* and *A. parasiticus*. Aflatoxin B1 (AFB1) is the most toxic and abundant aflatoxins (Jaynes and Zartman, 2011) contaminated in food or feeds in the field or during storage. AFB1 causes several toxicity as hepatotoxin, immunotoxin, carcinogenic and teratogenic to humans and animals (Jouany *et al.*, 2009; Mojtahedi *et al.*, 2013). Especially, AFB1 involved severe economic losses and health problems in dairy production because AFB1 is only partially degraded by rumen microorganisms. The remainder is then biologically converted to Aflatoxin M1 (AFM1) in the liver and eventually secreted in the dairy milk (Meucci *et al.*, 2011). The carry over rate of AFB1 in feed to AFM1 in milk were 0.01–0.02 (Jiang *et al.*, 2014). Moreover, previous studies reported that AFB1 decreased the gas production by affecting the rate and cumulative gas production (Jiang *et al.*, 2012; Mojtahedi *et al.* (2013). Aflatoxin also decreased cellulolysis in rumen (Edrington *et al.*, 1994).

As a result, various methods, including physical, chemical and biological method have been recently introduced to reduce or eliminate aflatoxin contaminated in feeds (Nemati *et al.*, 2015). Moreover, the mycotoxin adsorbents become the most commonly used to decontaminate mycotoxin in animal feed including clays, activated carbons and yeast. These products are capable of attaching mycotoxins (Mojtahedi *et al.*, 2013). Therefore, the objective of the study was to evaluate the effect of mycotoxin adsorbents for AFB1 decontamination and rumen fermentation efficiency by using *in vitro* gas production technique.

Materials and Methods

1. AFB1 Preparation

Commercially available AFB1 (Trilogy, Catalog number: TSL-401, USA) was prepared as stock standard solutions at 10 µg/ml using sterile deionized water.

2. Experimental design and treatments

The experimental design was conducted by using Completely Randomized Design (CRD). The basal diet containing 40% of rice straw and 60% of concentrate was allocated in eight treatments consisting T1=the control group, T2= the diet plus 1 µg/mg of aflatoxin B1 (AFB1), T3, T4, T5 = the diets supplement with 0.5% of each adsorbent (activated charcoal, montmorillonite, *S.cerevisiae*) and T6, T7, T8 = the diets plus 1 µg/mg of AFB1 together with 0.5% of each mycotoxin adsorbent, respectively. The dried basal diet was then milled, passed through a 0.2 mm sieve, and preceded to further step.

3. Rumen inoculums

Rumen fluid was collected from three healthy dairy cows, which were placed on a routine basis for at least 1 week and fed by the same basal diet for 5 days before sampling. *In vitro* fermentation was used in this study according to the technique described by Makkar *et al.* (1995). The rumen fluid (660 ml) was added to warm (39°C) and reduced medium consisting of 1,095 ml distilled water, 730 ml rumen buffer solution (417 mM NaHCO₃ and 51 mM NH₄HCO₃), 365 ml macromineral solution (46 mM KH₂PO₄, 40 mM Na₂HPO₄, 38 mM NaCl and 2 mM MgSO₄ · 7H₂O), 0.23 ml micromineral solution (505 mM MnCl₂ · 4H₂O, 898 mM CaCl₂ · 2H₂O, 42 mM CoCl₂ ·

6H₂O and 341 mM FeCl₂·6H₂O), 1 ml of 4mM resazurin and 60 ml freshly prepared reduction solution (145 mM Na₂S·9H₂O and 3.7 ml 1 M-NaOH). The mixture was kept stirring and continuously filled with CO₂ to ensure anaerobic condition at 39°C on a hot plate. Approximately 30 ml of the rumen-fluid medium was transferred into serum bottles incubated with feed samples (200 mg) at 39 °C for 72 h.

4. Sample collection and analysis

Dietary treatments were analyzed for dry matter (DM), ash, crude protein (CP) and ether extract (EE) using the procedure of AOAC (1997), while neutral detergent fiber (NDF) and acid detergent fiber (ADF) were also determined as described by Van Soest *et al.* (1991).

5. Gas production

During the incubation, the gas production kinetics were recorded at 1, 2, 4, 6, 8, 12, 24, 48 and 72 h. Cumulative gas production data were fitted to the model of Ørskov and McDonald (1979) as follow:

$$Y = a + b(1 - e^{-ct})$$

Where a = the gas production from the immediately soluble fraction, b = the gas production from the insoluble fraction, c = the gas production rate constant for the insoluble fraction (b), t = incubation time, (a+b) = the potential extent of gas production, y = gas production at time 't'. At 24- and 48-h post inoculation, a set of sample was determined *in vitro* true digestibility according to Van Soest and Robertson (1985).

6. Ammonia nitrogen and Volatile fatty acids analysis

Ruminal inoculum fluid was collected at 24 and 48 h after inoculation, about 30 ml of rumen inoculums mixed with 1M H₂SO₄ was centrifuged at 15,000xg for 15 min. Samples were divided into two portions for ammonia nitrogen (NH₃-N) analysis by using the micro-Kjeldahl methods (AOAC, 1990) and samples were then filtered through 0.22-µm nylon membrane (FILtrex, Singapore) for volatile fatty acids (VFAs) analysis by high performance liquid chromatography (HPLC) (model RF-10AXmugIL; Shimadze, Japan) according to Mathew *et al.* (1997).

7. Statistical analysis

All data were analyzed as a Completely Randomize Design using the SPSS version 19. The significant differences were compared by Post hoc test.

Results and Discussion

1. The chemical compositions

The chemical compositions of concentrate and rice straw in this study are shown in Table 1. The crude protein content in rice straw is 2.5%, 2.1 EE % and 83.0% OM, in accordance with previous report (Kang and Wanapat, 2013). Moreover, forage fiber analysis for rice straw in this experiment found that the rice straw contained 81.4% NDF, 50.8 % ADF and 5.9 ADL. The concentrate was formulated to contain 22.1% CP and 3.9% EE.

3. Gas production kinetic

Estimated parameters of gas production are presented in Table 2 and cumulative gas production profiles are shown in Fig. 1. In this

study, there were no treatment effect ($P>0.05$) on the fermentation of the soluble fraction (a) and effective gas production potential (EP). However, gas production from the insoluble fraction (b), gas production rate constant for the insoluble fraction and the potential extent of gas production were significantly different among treatments ($P<0.05$), of which the AFB1-containing diet was lower than the control group. These depressions in the gas production suggest that microbial populations are altered by AFB1 was add. This result was similar to previous studies (Jiang *et al.*, 2012; Mojtahedi *et al.*, 2013) where the asymptotic gas production numerically decreased by increasing AFB1 dosage.

The gas production from the insoluble fraction in the diet+AFB1 was lower than in the diet + AFB1 + montmorillonite and diet + AFB1 + *S. cerevisiae*. In addition, the gas production rate was the lowest in treatment plus AFB1. However, montmorillonite and *S. cerevisiae* were effectively decreased the AFB1 effect on gas production. This phenomenon could be explained through the detoxifying capability of *S. cerevisiae* cell wall and montmorillonite. The adsorption capacity largely

depended on both yeast composition and mycotoxin as well. Joannis-Cassan *et al.* (2011) reported that the yeast cell wall from baker's yeast could absorb up to 29% of AFB1. Similarly, Desheng *et al.* (2005) also noted that the montmorillonite absorbed more than 80% of the available AFB1 from aqueous solution at pH 2 and 90% at pH 8, correspondingly previous report, where montmorillonite could absorb up to 93% of AFB1 (Dakovi *et al.*, 2008).

4. Degradability and rumen metabolites

The *in vitro* dry matter digestibility (IVDMD) and organic matter digestibility (IVOMD) were not significantly affected by AFB1 and mycotoxin adsorbents ($P>0.05$) (Table 3). Likewise, Pettersson and Kiessling (1976) and Jiang *et al.* (2012) demonstrated that addition of AFB1 did not alter IVDMD. However, the result was opposed with Mojtahedi *et al.* (2013) who reported that IVDMD decreased significantly with AFB1 inclusion. It could be explained that AFB1 could attributed to compromise ruminal fermentation by reducing fiber digestion and VFA production.

Table 1 The chemical composition of rice straw and concentrate

Items	Rice straw	Concentrate
DM, %	94.3	91.9
Chemical composition	-----% of DM basis-----	
OM	83.0	83.2
Ash	12.5	7.8
CP	2.5	22.1
EE	2.1	3.9
NDF	81.4	21.1
ADF	50.8	11.7
ADL	5.9	3.6

DM = Dry matter, OM = Organic matter, CP = Crude protein, EE = Ether extract
 NDF = Neutral- detergent fiber, ADF = Acid- detergent fiber, ADL = Acid- detergent lignin

Table 2 Effects of AFB1 and mycotoxin adsorbents on *in vitro* gas production of dietary treatments

Items	Treatments								SEM	P-value
	T1	T2	T3	T4	T5	T6	T7	T8		
Gas production characteristics										
a	1.21	1.53	-0.64	-0.25	-0.48	-2.16	-1.55	-0.97	0.36	0.13
b	50.7 ^{bcd}	38.7 ^a	53.7 ^{cde}	44.2 ^{ab}	58.6 ^e	50.9 ^{abc}	56.2 ^{de}	49.5 ^{bcd}	1.54	<0.01
c	0.07 ^{ab}	0.18 ^c	0.08 ^{ab}	0.13 ^c	0.07 ^{ab}	0.12 ^{bc}	0.06 ^a	0.11 ^{abc}	0.01	0.02
d	51.9 ^{bc}	40.2 ^a	53.0 ^{bc}	44.0 ^{ab}	58.1 ^c	48.7 ^{abc}	54.6 ^{bc}	48.5 ^{abc}	1.52	0.04
Effective gas production potential (EP)										
	31.9	30.4	31.5	31.3	33.2	33.2	30.1	32.2	0.53	0.82

^{a,b,c,d,e} Means in the same row with different superscript differ (P<0.05)

a = the gas production from the immediately soluble fraction, b = the gas production from the insoluble fraction, c = the gas production rate constant for the insoluble fraction, d = the potential extent of gas production, EP = a + [bc/(k + c)] where k = 0.05 (Ørskov and McDonald, 1979).

T1=Basal diet (the control), T2= Basal diet plus AFB1 at 1 µg/mg, T3= Basal diet supplement with 0.5% activated charcoal,

T4= Basal diet supplement with 0.5% montmorillonite, T5= Basal diet supplement with 0.5% *S.cerevisiae*,

T6= Basal diet plus AFB1 at 1 µg/mg supplement with 0.5% activated charcoal,

T7= Basal diet plus AFB1 at 1 µg/mg supplement with 0.5% montmorillonite,

T8= Basal diet plus AFB1 at 1 µg/mg supplement with 0.5% *S. cerevisiae*




Fig. 1 Effect of AFB1 and mycotoxin adsorbents on gas production at different times of incubation in dairy cow's rumen fluid

T1=Basal diet (the control), T2= Basal diet plus AFB1 at 1 µg/mg, T3= Basal diet supplement with 0.5% activated charcoal,

T4= Basal diet supplement with 0.5% montmorillonite, T5= Basal diet supplement with 0.5% *S.cerevisiae*,

T6= Basal diet plus AFB1 at 1 µg/mg supplement with 0.5% activated charcoal, T7= Basal diet plus AFB1 at 1 µg/mg supplement

with 0.5% montmorillonite, T8= Basal diet plus AFB1 at 1 µg/mg supplement with 0.5% *S. cerevisiae*

The results of VFAs (Acetic acid, Propionic acid, and Butyric acid) and NH₃-N are shown in Table 4. In this study, there were no effects of treatments on the VFAs and NH₃-N (P>0.05). However, some studies reported that the addition of AFB1 *in vitro* rumen fermentation significantly decreased NH₃-N concentrations (Mojtahedi *et al.*

2013; Jiang *et al.*, 2012). AFB1 maybe was affected ammonia. Protein synthesis and increased the microbial requirement for nutrients (Edrington *et al.*, 1994) when the digestion of protein was inhibited by AFB1, resulting in the decreased ammonia concentration.

Table 3 Effects of AFB1 and mycotoxin adsorbents on *in vitro* digestibility

Items	Treatments								SEM	P-value
	T1	T2	T3	T4	T5	T6	T7	T8		
<i>In vitro</i> dry matter digestibility (IVDMD), %										
24 h	48.9	49.5	49.5	47.5	50.7	48.9	53.2	52.2	0.64	0.42
48 h	58.9	59.2	58.0	62.1	56.2	61.8	57.7	62.2	0.38	0.55
<i>In vitro</i> organic matter digestibility (IVOMD), %										
24 h	54.8	54.2	55.3	53.8	56.5	56.1	58.0	56.8	0.52	0.58
48 h	65.4	65.2	63.7	66.9	63.2	66.1	64.1	66.1	0.63	0.89

T1=Basal diet (the control), T2= Basal diet plus AFB1 at 1 µg/mg, T3= Basal diet supplement with 0.5% activated charcoal, T4= Basal diet supplement with 0.5% montmorillonite, T5= Basal diet supplement with 0.5% *S.cerevisiae*, T6= Basal diet plus AFB1 at 1 µg/mg supplement with 0.5% activated charcoal, T7= Basal diet plus AFB1 at 1 µg/mg supplement with 0.5% montmorillonite, T8= Basal diet plus AFB1 at 1 µg/mg supplement with 0.5% *S. cerevisiae*

Table 4 Effects of AFB1 and mycotoxin adsorbents on volatile fatty acids and ammonia nitrogen

Items	Treatments								SEM	P-value
	T1	T2	T3	T4	T5	T6	T7	T8		
Total VFAs, mM										
24 h	112.7	108.1	109.1	125.9	111.5	91.3	122.3	126.3	3.28	0.07
48 h	133.5	122.6	127.8	137.2	120.2	128.1	131.5	127.8	2.40	0.80
Acetic acid (C ₂), %										
24 h	58.9	55.1	56.1	58.1	56.5	53.0	58.0	58.2	0.67	0.41
48 h	60.5	58.1	59.8	59.8	58.9	59.2	59.4	59.3	0.37	0.92
Propionic acid (C ₃), %										
24 h	25.7	32.4	30.5	30.0	29.5	33.1	27.8	27.7	0.79	0.27
48 h	27.1	27.9	27.9	28.2	28.9	28.0	27.2	27.2	0.28	0.84
Butyric acid (C ₄), %										
24 h	15.5	12.5	13.5	13.0	14.0	14.0	14.3	14.1	0.27	0.17
48 h	12.4	14.1	12.2	12.0	12.2	12.8	13.3	13.6	0.22	0.10
C ₂ :C ₃ ratio										
24 h	2.37	1.72	1.86	2.01	1.91	1.60	2.09	2.10	0.08	0.38
48 h	2.23	2.08	2.15	2.12	2.05	2.12	2.19	2.18	0.03	0.94
Ammonia nitrogen (NH ₃ -N), mg%										
24 h	17.6	17.0	18.2	15.2	15.2	18.2	17.0	18.0	0.29	0.39
48 h	18.9	29.1	18.8	18.5	19.5	20.1	21.3	21.6	0.36	0.20

T1=Basal diet (the control), T2= Basal diet plus AFB1 at 1 µg/mg, T3= Basal diet supplement with 0.5% activated charcoal, T4= Basal diet supplement with 0.5% montmorillonite, T5= Basal diet supplement with 0.5% *S.cerevisiae*, T6= Basal diet plus AFB1 at 1 µg/mg supplement with 0.5% activated charcoal, T7= Basal diet plus AFB1 at 1 µg/mg supplement with 0.5% montmorillonite, T8= Basal diet plus AFB1 at 1 µg/mg supplement with 0.5% *S. cerevisiae*

Conclusion

In conclusion, our study found that AFB1 at 1 µg/mg affected *in vitro* fermentation characteristics by reducing the gas production from the insoluble fraction and potential gas production. However, the gas production from the immediately soluble fraction, EP, NH₃-N and VFAs concentrations did not differ among treatments. In addition, montmorillonite and *S. cerevisiae* effectively reduced the effects of AFB1 on gas production.

Acknowledgements

This research was supported by Faculty of Veterinary Medicine, Khon Kaen University and Higher Education Research Promotion and National Research University Project of Thailand, Office of the Higher Education Commission, the Food and Functional Food Research of Khon Kaen University for financially support this research project.

References

- AOAC. 1990. Official methods of analyses, 15th Ed., Association of Official Agriculture Chemicals. Arlington, VA.
- AOAC. 1997. Official Method of Analysis. 16th Ed., Association of Official Agriculture Chemicals. Washington, DC. USA.
- Dakovic, A., Matijasevic, S., Rottinghaus, G.E., Ledoux, D.R., Butkeraitis, P. and Sekulic, Z. 2008. Aflatoxin B1 adsorption by natural and copper modified montmorillonite. *Biointerfaces*. 66: 20-25.
- Desheng, Q., Fan, L., Yanhu, Y. and Niya, Z. 2005. Adsorption of Aflatoxin B1 on Montmorillonite. *Poult. Sci.* 84: 959-961.
- Edrington, T.S., Harvey, R.B. and Kubena, L.F. 1994. Effect of aflatoxin in growing lambs fed ruminally degradable or escape protein sources. *J. Anim. Sci.* 72: 1274-1281.
- Jaynes, W.F. and Zartman, R.E. 2011. Aflatoxin toxicity reduction in feed by enhanced binding to surface modified clay additives. *Toxins*. 3: 551-565.
- Jiang, Y.H., Wang, P., Yang, H.J. and Chen, Y. 2014. The Efficacy of bamboo charcoal in comparison with smectite to reduce the detrimental effect of aflatoxin B1 on *in vitro* rumen fermentation of a hay-rich feed mixture. *Toxins*. 6: 2008-2023.
- Jiang, Y.H., Yang, H.J. and Lund, P. 2012. Effect of aflatoxin B1 on *in vitro* ruminal fermentation of rations high in alfalfa hay or ryegrass hay. *J. Feed Sci. Technol.* 175: 85-89.
- Joannis-Cassan, C., Tozlovanu, M., Hadjeba-Medjdoub, H., Ballet, N. and Leszkowicz, A.P. 2011. Binding of zearalenone, aflatoxin B1, and ochratoxin A by yeast-based products: a method for quantification of adsorption performance. *J. Food. Prot.* 2011: 1175-1185.
- Jouany, I.P., Yiannikouris, A. and Bertin, G. 2009. Risk assessment of mycotoxins in ruminants and ruminant production. *Nutri. foraging ecology sheep and goats*. 85: 205-224.

- Kang, S. and Wanapat, M. 2013. Using plant source as a buffering agent to manipulating rumen fermentation in an *in vitro* gas production system. *Asian-Austra. J. Anim. Sci.* 26: 1424-1436.
- Makkar, H.P.S., Blummel, M. and Becker, K. 1995. Formation of complexes between polyvinyl pyrrolidones or polyethylene glycols and tannins, and their implication in gas production and true digestibility in *in vitro* techniques. *Br. J. Nutr.* 73: 897-913.
- Mathew, S., Sagathevan, S., Thomas, J. and Mathen, G. 1997. An HPLC method for estimation of volatile fatty acids in ruminal fluid. *Indian. J. Anim. Sci.* 67: 805-807.
- Meucci, V., Soldani, G., Razzuoli, E., Saggese, G. and Massart, F. 2011. Mycoestrogen pollution of Italian infant food. *J. pediatr.* 152: 278-283.
- Mojtahedi, M., Danesh, M., Vakili, S.A. and Hayati-Ashtiani, M. 2013. Effect of aflatoxin B1 on *in vitro* rumen microbial fermentation responses using batch culture. *Annual rev research in biology.* 3: 686-693.
- Nemati, Z., Karimi, A. and Besharati, M. 2015. Effects of aflatoxin B1 and yeast cell wall supplementation on the growth performance of broilers. *International Conference on Innovations in Chemical and Agricultural Engineering.* Kuala Lumpur: Malaysia.
- Ørskov, E.R. and McDonald, I. 1979. The estimation of protein degradability in the rumen from incubation measurements weighted according to rate of passage. *J. Agri. Sci.* 92: 499 -503.
- Pettersson, H. and Kiessling, K.H. 1976. Metabolism of aflatoxin by intact rumen fluid, rumen protozoa and rumen bacteria. *J. Agric Res.* 6:161.
- Van Soest, P.J. and Robertson, J.B. 1985. *Analysis of Forages and Fibrous Foods. A Laboratory Manual for Animal Science.* Department of Animal Science, Cornell University, Ithaca, NY: USA.
- Van Soest, P.J., Robertson, J.B. and Lewis, B.A. 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74: 3583-3597.