



Contamination of Microplastics in Retail *Paratapes undulatus* Clams from Fresh Markets in Nakhon Pathom Province, Thailand

Pattrawan Khamboonruang, Mint Rueawraengbunya & Taeng On Prommi*

Faculty of Liberal Arts and Science, Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom Province, 73140 Thailand

Article info

Article history:

Received: 16 November 2023

Revised: 16 December 2023

Accepted: 19 December 2023

Keywords:

Microplastics, *Paratapes undulatus*, FTIR, Human health

Abstract

Microplastic contamination in food is a growing problem in the modern world. Clams, in particular, are consumed whole and are particularly susceptible to contamination entering the body. In 120 clams, microplastic contamination was investigated. Clam tissues were digested with 30% hydrogen peroxide and 10% potassium hydroxide. A total of 1,001 microplastic items were found. There was a statistically significant difference between the weights of clam tissue and microplastic ($\chi^2 = 36.945$, $df = 5$, $p = 0.000$; $\chi^2 = 35.842$, $df = 5$, $p = 0.000$). The most prevalent microplastic shapes, at 38% and 36%, were identified as fragment and film microplastics. The most common microplastic color was white or transparent (29%), whereas the majority (44%) of the microplastics were less than 100 μm . A Fourier transform infrared spectrometer (FTIR) was used to confirm 99 microplastics that were chosen at random and amounted to approximately 10% of the total microplastics. PET (polyethylene terephthalate), BEHP (bis(2-ethylhexyl) phthalate), PP (polypropylene), CA (cellulose acetate), PVAc (polyvinyl acetate), PMMA (poly (methyl methacrylate) and EC (ethyl cellulose) are examples of plastic materials. According to the study's findings, the *Paratapes undulatus* sold in Thailand's fresh market was heavily contaminated with microplastics. As a result, the findings of this study can be utilized to inform future research on assessing exposure to microplastics and the health risks associated with consuming contaminated bivalves commonly consumed in Thailand.

Introduction

Microplastics (plastic particles smaller than 5 mm) (Frias & Nash, 2019) have been identified as one of the most concerning contaminants in marine and

coastal habitats worldwide (Harding, 2016; Shahul Hamid et al., 2018). According to reports, between 4.8 and 12.7 million metric tons of plastic trash from 192 coastal nations had entered marine habitats by 2010 (Jambeck et al., 2015), making plastic debris the most

* Corresponding Author
e-mail: faastop@ku.ac.th

prevalent type of marine litter, accounting for three-quarters of marine waste (Harding, 2016). Worryingly, this number is expected to rise substantially in the coming years, with plastic items expected to reach 33 billion tons by 2050 (Harding, 2016). Because of the persistence and abundance of plastic waste in the ecosystem, larger plastics have broken down into secondary microplastics, resulting in a huge number of microplastics in all coastal and marine environments, including estuaries, mangroves, lagoons, bays and deep-sea areas (Hitchcock & Mitrovic, 2019; Nor & Obbard, 2014; Bayo et al., 2019; Falahudin et al., 2020; Van Cauwenberghe et al., 2013). Additionally, due to product discharges containing microplastics after use or accidents in plastic transportation, primary microplastics-plastics made for a specific purpose (e.g., plastic pellets for drugs and cosmetics products)-are a major source of pollution in the environment (Pandey et al., 2022). Microplastics have been spread to all levels of trophic organisms in the food web due to their extensive availability in all coastal and marine ecosystems. According to Gall & Thomson (2015), plastic accounts for 92% of all contacts between individual creatures and marine litter, affecting these species through ingestion, entanglement and habitat disruption. The major effects of microplastics on aquatic habitats include physical damage, atypical behavior patterns, obstacles in the nutrient cycle, cytotoxicity and genotoxicity and a rise in mortality (Pandey et al., 2022).

The bivalve mollusks are filter feeders and belong to species that are most vulnerable to microplastic contamination due to their feeding habits, as they ingest suspended particles from the water column (Setälä et al., 2016; Ward et al., 2019; Li et al., 2019). These microplastics can potentially have a wide range of deleterious impacts on bivalves, including affecting filtering function and reproduction systems, causing genotoxicity and producing indirect consequences related to modifying the structure of habitat and food supplies (Zhang et al., 2020). In Asian green mussels (*Perna viridis*), exposure to polyvinyl chloride (PVC) raised mortality rates, whereas in Pacific oysters (*Crassostrea gigas*), exposure to polystyrene (PS) affected sperm motility, egg production and oocyte quality (Pandey et al., 2022). Various organisms, including invertebrates, fish, birds and mammals, rely on bivalves for food (Waser, 2018). As a result, these faunas can move microplastics up the food chain to higher-trophic species. More significantly, bivalves are a significant source of sustenance for humans; their yearly production exceeds 15 million tons,

or 14% of all marine production worldwide (Smaal, 2019). Because humans consume all of the tissues of bivalves, unlike other seafood species, bivalves may be a significant source of microplastic for humans (Zhang et al., 2020). Microplastics have been found in a variety of economically popular bivalves, such as clams, scallops, oysters and mussels, throughout natural and aquaculture habitats (Qu et al., 2018; Naji et al., 2018; Phuong et al., 2018; Davidson et al., 2016; Li et al., 2016; Van Cauwenberghe & Janssen, 2014). Microplastic concentration in bivalves is commonly between 10^{-1} and 10^1 items g^{-1} (Cho et al., 2019; Jin-Feng et al., 2018; Renzi et al., 2018). In Canadian bivalve species, however, microplastic amounts up to 657.5 items g^{-1} have been reported (Murphy, 2018). Van Cauwenberghe & Janssen (2014) estimate that Europeans with an average mollusk intake of 11.8-72.1 g $cap^{-1} day^{-1}$ may consume up to 1,800-11,000 microplastics per year, which can have serious consequences for human health.

Bivalve mollusks are a popular type of seafood in Thailand, among locals and tourists. The government intends clams, in particular, to be one of Thailand's main seafood exports to nations worldwide. However, data on the level of microplastic contamination in edible bivalves is currently scarce, despite the fact that microplastics have been identified in quite significant concentrations across the country throughout freshwater systems and coastal areas (Strady et al., 2021; Tran-Nguyen et al., 2020). As a result, microplastics are highly likely to accumulate in living organisms and negatively affect consumer health. The presence and characteristics of microplastics in the clam *Paratapes undulatus*, a common edible bivalve mollusk in Thailand, were investigated in this study.

A wide range of acids (Van Cauwenberghe & Janssen, 2014), enzymes (Catarino et al., 2017), alkalis (Rochman et al., 2015) and oxidizing agents, such as H_2O_2 (Li et al., 2015), are frequently used in the digestion of bivalve tissue. Numerous steps in certain digestion procedures raise the risk of airborne contaminants getting into the sample. As a result, this study determined the most appropriate approach for breaking down bivalve tissues and extracting microplastics by contrasting the KOH and H_2O_2 approaches used in previous research. The study describes a method for the extraction and quantification of microplastics from clams. This study will add to the understanding of microplastic contamination in Thai biota as well as the implications for humans, including dietary exposure, in order to determine the potential threat of contaminated seafood.

Materials and methods

1. Sample collection and preparation

In the Gulf of Thailand and the Andaman Sea, the clam *P. undulatus* (Born, 1778) supports Thailand's largest shellfishery. In general, the harvest areas for clams are consolidated mud substrates within 3 to 7 kilometers of the shoreline. Shellfish grounds can be found off the coasts of various coastal provinces and are collected all year. Clams are typically acquired for export as canned products. A total of 120 individuals of *P. undulatus* (approximately 1 kilogram) were purchased at the fresh market in Nakhon Pathom Province for an investigation of microplastics (MPs). These clams were harvested from Samut Prakarn Province. All *P. undulatus* samples were rinsed with filtered distilled water. The shell lengths of *P. undulatus* were also measured. *P. undulatus* samples were separated from their shells prior to tissue processing and the soft tissue of each sample was weighed separately. The tissue from each sample was preserved in the freezer until it was time to be tested.

In this investigation, the clam tissue digestion studies were separated into two groups: H₂O₂ and KOH. Each group had three replications, each of which had 20 individual clams.

2. Microplastic isolation from the soft tissue of *P. undulatus*

Clam tissue digestion was carried out, according to Ehlers et al. (2019). To break down the soft tissue, each individual was put down in a labeled 100-mL Erlenmeyer flask that had already been cleaned and approximately 20 mL of 30% H₂O₂ was poured into each flask. Each flask was wrapped in parafilm. It was then boiled at 60°C for 3 hr in a shaken water bath at 150 rpm, or until all of the organic stuff was digested. The blanks were checked in parallel for the presence of MPs. In the other group, 20 mL of 10% KOH was introduced to each flask containing clam soft tissue at the same time.

Microplastics (MPs) were recovered from a dissolved organic matter solution via flotation with potassium formate (HCO₂K) (Zhang et al., 2016). 99% HCO₂K was added to each sample in a glass separating funnel until the concentration reached 1.6 g/mL. The samples were then kept at room temperature for a minimum of 3 hr. A layer of MPs emerged when the less dense particles separated from the saturated solution, resulting in the undissolved organic residues and inorganic materials sinking to the bottom of the glass containers. Subsequently, the samples were filtered using a pressure filtering system

and a nylon membrane filter (Whatman, Kent, UK; pore size, 0.45 µm; diameter, 47 mm). After being filtered, each membrane was put on a sterile Petri dish, covered with aluminum foil and allowed to dry for two days at 50°C in a drying cabinet.

3. Microscopic examination of MPs

The prevalence and characteristics of MPs in filtered filters were determined using a stereomicroscope equipped with a camera and image analysis software (Leica EZ4E, Germany). The microplastic particle size was measured using the LAS X software and the length was calculated from the longest side. A visual inspection was also done to identify expected MPs based on morphological characteristics such as color and shape (Hidalgo-Ruz et al., 2012). The kinds of microplastic particles were classified as fiber, spherical, film (a thin and small layer) and fragment (part of a larger plastic item) (Su et al., 2016; 2018). MPs were divided into four groups, with *L* representing the length of the longest diameter: first ($L \leq 100 \mu\text{m}$); second ($100 < L \leq 250 \mu\text{m}$); third ($250 < L \leq 500 \mu\text{m}$); and fourth ($L > 500 \mu\text{m}$).

4. Polymer type identification

To determine the polymer types, a representative number of MPs from each morphotype were randomly picked and evaluated with a PerkinElmer Spectrum-Fourier transform infrared spectrometer (FTIR) in attenuated total reflection (ATR) mode. The MPs chosen reflected the most common types of visually observed particles in all samples. A Hyperion 2000 FTIR microscope (Bruker Daltonik, Billerica, MA, USA) with a mercury-cadmium telluride detector was used to manually analyze 99 particles from soft tissue at wave numbers ranging from 4000 to 600 cm⁻¹, with 32 co-added scans and a spectrum resolution of 4 cm⁻¹. The Bruker spectrum library was used to compare polymer types and functional group characterizations. Based on the spectrum analysis, a matching level of spectra with a quality index of ≥ 0.7 was considered acceptable (Woodall et al., 2014).

5. Quality control

To prevent airborne microplastic contamination, non-plastic equipment such as cotton lab coats and nitrile or latex gloves was worn throughout the experimental procedures. In addition, glassware and metal tools were utilized in the lab to evaluate the samples. During filtering and sieving, any potential contaminants were discovered. Air exposure was kept to a minimum in order to reduce microplastic loss.

6. Statistical analysis

Microplastics in clam tissue were measured

individually based on type, color and size. Pooled samples (20 specimens, 6 replicates) were used to calculate the average amount of MPs per g of wet weight and individual clam tissue. Furthermore, for each replication, the Chi-square test of independence was used to assess microplastic shape, color and size. All statistical tests were run on Statistica 20.0, a computer program.

Results and Discussion

1. Abundance of microplastics

The shell length of the 120 individual *Paratapes undulatus* was 4.32 ± 0.38 cm and the wet tissue weight was 1.82 ± 0.44 g. Microplastic contamination in all *P. undulatus* was detected in a total of 1,001 suspected microplastic items (Table 1). *P. undulatus* had a microplastic abundance of 4.88 ± 1.65 items g^{-1} ($8.34 \pm$

3.87 items individual^{-1}). Our results confirmed the presence of MPs in all of the clams analyzed.

In experiment H_2O_2_2 , the average microplastics were detected at 13.70 ± 12.25 items individual^{-1} (7.61 ± 5.61 items g^{-1}), followed by experiment H_2O_2_3 , where microplastics were found at 12.90 ± 5.60 items individual^{-1} (8.41 ± 4.56 items g^{-1}), KOH_1 discovered 7.00 ± 5.71 items individual^{-1} (3.85 ± 3.34 items g^{-1}), H_2O_2_1 detected 6.15 ± 7.90 items individual^{-1} (3.18 ± 3.99 items g^{-1}), KOH_2 found 5.15 ± 1.90 items individual^{-1} (3.20 ± 1.56 items g^{-1}) and KOH_3 found 5.15 ± 2.08 items individual^{-1} (3.02 ± 1.48 items g^{-1}) (Fig. 1, Table 1). The weights of clam tissue and microplastic digested by hydrogen peroxide and potassium hydroxide differed statistically ($\chi^2 = 36.945$, $df = 5$, $p = 0.000$; $\chi^2 = 35.842$, $df = 5$, $p = 0.000$) (Fig. 1).

Table 1 The weight of clam flesh and the number of microplastics (MPs) found after digesting clam tissue with hydrogen peroxide and potassium hydroxide

30% H_2O_2_1			30% H_2O_2_2			30% H_2O_2_3			10% KOH_1			10% KOH_2			10% KOH_3		
No.	Edible tissue (g)	MPs (item)	No.	Edible tissue (g)	MPs (item)	No.	Edible tissue (g)	MPs (item)	No.	Edible tissue (g)	MPs (item)	No.	Edible tissue (g)	MPs (item)	No.	Edible tissue (g)	MPs (item)
1	2.01	30	1	1.61	24	1	1.97	18	1	2.94	6	1	1.74	5	1	1.99	6
2	1.97	9	2	2.46	41	2	1.49	15	2	1.93	7	2	1.23	7	2	1.90	6
3	1.56	3	3	1.40	13	3	1.13	12	3	1.36	7	3	2.13	6	3	1.32	9
4	1.80	16	4	1.10	12	4	1.06	6	4	1.82	8	4	3.50	1	4	1.60	2
5	2.19	19	5	1.76	7	5	1.51	14	5	1.72	28	5	1.56	9	5	2.03	6
6	2.16	5	6	1.72	3	6	0.96	15	6	1.48	7	6	1.87	6	6	1.69	2
7	2.11	1	7	2.05	8	7	1.30	11	7	2.15	7	7	1.28	7	7	1.47	4
8	1.53	3	8	1.50	6	8	1.89	13	8	1.91	11	8	2.16	4	8	1.55	5
9	2.56	1	9	2.10	5	9	1.45	20	9	2.72	9	9	1.63	5	9	1.75	2
10	1.45	4	10	2.70	44	10	1.99	15	10	1.69	2	10	1.51	6	10	1.85	4
11	2.14	2	11	1.70	7	11	1.38	29	11	1.81	12	11	1.36	7	11	2.67	6
12	2.35	1	12	1.63	5	12	2.05	13	12	1.87	6	12	2.07	4	12	1.37	4
13	2.04	10	13	1.71	33	13	1.75	12	13	1.78	7	13	1.94	7	13	1.48	6
14	2.65	1	14	1.74	9	14	2.87	7	14	1.39	5	14	1.56	5	14	1.51	7
15	1.91	1	15	1.77	3	15	1.72	5	15	2.10	5	15	1.73	2	15	1.68	10
16	3.12	2	16	1.70	17	16	1.44	16	16	1.20	3	16	1.31	5	16	1.86	4
17	2.14	1	17	1.53	7	17	2.56	10	17	1.99	3	17	1.90	3	17	2.26	6
18	1.70	12	18	2.26	9	18	1.98	13	18	1.79	3	18	1.61	4	18	2.36	4
19	1.74	1	19	1.74	7	19	1.32	9	19	2.35	2	19	1.38	6	19	1.34	5
20	1.26	1	20	1.14	14	20	1.59	5	20	2.15	2	20	2.35	4	20	1.82	5
sum	40.38	123		35.32	274		33.41	258		38.13	140		35.80	103		35.46	103
mean	2.02	6.15		1.77	13.70		1.67	12.90		1.91	7.00		1.79	5.15		1.77	5.15
SD	0.44	7.90		0.39	12.25		0.48	5.60		0.43	5.71		0.51	1.90		0.36	2.08

Remark: H_2O_2_1 = digesting clam tissue with hydrogen peroxide in the first replication; H_2O_2_2 = digesting clam tissue with hydrogen peroxide in the second replication; H_2O_2_3 = digesting clam tissue with hydrogen peroxide in the third replication; KOH_1 = digesting clam tissue with potassium hydroxide in the first replication; KOH_2 = digesting clam tissue with potassium hydroxide in the second replication; KOH_3 = digesting clam tissue with potassium hydroxide in the third replication

The average amount of microplastics ingested by clams in this study (8.34 ± 3.87 items individual⁻¹ or 4.88 ± 1.65 items g⁻¹) is higher than values reported in coastal waters of China (1.5 to 7.6 items individual⁻¹) (Li et al., 2016) and the UK (1.1–6.4 items individual⁻¹) (Li et al., 2018), as well as in Giglio Island, Italy (1–2 items individual⁻¹) (Avio et al., 2017) and the French Atlantic coast (0.6 ± 0.6 items individual⁻¹) (Phuong et al., 2018). According to the reports studied, the Gulf of Thailand has significant microplastic contamination (Imasha & Babel, 2023; Srikrajang & Prommi, 2021). However, of the total debris items reported in the UK mussel study (Li et al., 2018), only 50% were microplastics. The microplastic ingestion values of our study are much higher than those of Vandermeersch et al. (2015) in the estuaries of Portugal, Italy and Spain; De Witte et al. (2014) in the Belgian coast; and Van Cauwenberghe et al. (2015) in the French, Belgian and Dutch North Sea coasts. The previous three studies provided results per weight of complete mussel tissues, which may explain the observed differences from our results, which are calculated per weight of gills and digestive tract and include higher microplastic quantities among mussel tissues (Tsangaris et al., 2015). Moreover, previous investigations utilized pooled samples and reported mean values for all individuals studied, whereas we computed the number of microplastics on a wet-weight basis, considering only those with microplastics. Our findings are not comparable to other studies that show microplastic ingestion in mussels on a dry weight basis (Leslie et al., 2017; Karlsson et al., 2017). Most mussel studies do not reveal the frequency of ingested microplastics, which is likely due to the use of pooled animal samples for microplastic extraction. Microplastic occurrence rates in mussels in our study (100%) are higher than those reported by Avio et al. (2017) in Giglio Island, Italy (10–36%), who employed individual mussels for microplastic extraction in a similar manner. The variations that were observed could be related to the use of different mussel sampling strategies; for example, in our research, we collected both wild and cultured clams, whereas Avio et al. (2017) implanted mussels at two depths (surface and bottom) for four weeks.

2. Microplastic shape, size, and color

On average, each sample contained eight microplastic items. In the present study, the most prevalent microplastics were fragments, followed by films and fibers. Approximately 38% of the microplastics obtained from *P. undulatus* soft tissues were fragment-shaped.




Fig. 1 Mean concentrations of microplastics found in *Paratapes undulatus* clams after digesting clam tissue with hydrogen peroxide and potassium hydroxide

Microplastics in the film shape accounted for 36% of the total, whereas microplastics in the fiber shape accounted for 26% of the total (Fig. 2a). The microplastic shape in clam tissue degraded by hydrogen peroxide and potassium hydroxide differed statistically in fragment, fiber and film ($\chi^2 = 25.126$, $df = 5$, $p = 0.000$; $\chi^2 = 12.739$, $df = 5$, $p = 0.000$; $\chi^2 = 70.222$, $df = 5$, $p = 0.000$, respectively) (Fig. 3). Mechanical and UV radiation are likely to break down fragments from larger plastic objects (e.g., plastic containers, furniture and toys) (Horton et al., 2017). Thus, this study's higher percentage of fragments indicates that MPs found in the soft tissue are mainly secondary microplastics. In this investigation, the composition of ingested microplastics in market clams revealed higher levels of fragments than fibers and film. Similarly, Phuong et al. (2018) found that most microplastics in mussels from the French Atlantic coast were fragments, reaching 82%, greater than our result (38% fragments). However, most mussel investigations (De Witte et al., 2014; Li et al., 2016) show that fibers predominate fragments. Different sources and waste management practices could explain the variation in shape categories of ingested microplastics among market clams and other places (Rochman et al., 2015). Poor waste management methods, both on land and on the shoreline, where tourism and leisure activities are intense, are among the litter inputs in the Gulf of Thailand, where the clams were harvested. This may result in large amounts of plastic inputs into the sea (e.g., plastic bags, bottles and cups) that can degrade into microplastic fragments. Furthermore, poorly managed plastic can fragment more quickly on land and enter the sea as microplastics (Kalogerakis et al., 2017). Plastic particles were found in similar proportions in ingested microplastics across species, habitats and sampling sites, indicating widespread distribution in the study area.




Fig. 2 Composition of microplastic (MPs) found in clam tissue; a) morphotype b) size and c) color




Fig. 3 Mean concentrations of microplastic shapes found in *Paratapes undulatus* after digesting clam tissue with hydrogen peroxide and potassium hydroxide

Microplastics were divided into four sizes based on the largest dimension: < 100 μm , 100–250 μm , 250–500 μm , > 500 μm . Microplastic particles with a diameter of less than 100 μm (44%) were the most abundant, followed by those with a diameter of 100–250 μm (29%) and 250–500 μm (17%). Particles larger than 500 μm (10%) were less common (Fig. 2b). The microplastic size in clam tissue degraded by hydrogen peroxide and potassium hydroxide found that all sizes

(<100 μm , 100–200 μm , 200–500 μm and >500 μm) differed statistically ($\chi^2 = 53.501$, $df = 5$, $p = 0.000$; $\chi^2 = 27.885$, $df = 5$, $p = 0.000$; $\chi^2 = 16.701$, $df = 5$, $p = 0.000$; $\chi^2 = 19.203$, $df = 5$, $p = 0.000$) (Fig. 4). In terms of microplastic size, particles less than 100 μm were the most common, followed by particles 100–250 μm . Microplastics with diameters smaller than 0.1 mm, on the other hand, may have been underestimated, as recovery rates decline with particle size (Avio et al., 2015). Suaria et al. (2016) found that the existence of particles with a diameter of less than 0.5 mm conforms to the size distribution of floating microplastics in Mediterranean waterways. Because of their filter-feeding approach, the clams ingested more microplastics of smaller sizes (less than 0.1 mm, 0.1 mm–0.5 mm).




Fig. 4 Mean concentrations of microplastic sizes found in *Paratapes undulatus* after digesting clam tissue with hydrogen peroxide and potassium hydroxide

According to Figs. 2a and 4, the size ranges of less than 100 μm and 100–250 μm were the most common classes for *P. undulatus*, representing 44% and 29%, respectively. Microplastics that are larger than 500 μm exhibit 10% of *P. undulatus*. The feeding mode is hypothesized to influence the size range of digested microplastics. Small-size microplastics are a significant component of marine animal ingestion because they make plastic particles extensively accessible to various biota in both benthic and pelagic environments. Smaller particles appear to be more likely to be swallowed, resulting in stomach blockage, physiological consequences, chemical transfer and trophic transfer, whereas larger particles are more likely to entangle, preventing marine animals from swimming, filtering, or catching their prey (Cole et al., 2011; Lehtiniemi et al., 2018; Lusher, 2015). Some feeding processes do not allow them to differentiate between prey and anthropogenic materials in some situations, while others may mistake it for food and consume the plastic directly (Lusher, 2015).

Twelve distinct hues of microplastic particles were found (Fig. 2c): white, black, transparent, light blue, navy blue, green, silver, brown, yellow, red, pink and violet. The most prevalent colors of microplastics in clam samples were white (28%), black (16%), navy blue (15%) and brown (10%); some were yellow (6%), light blue (3%), red (3%), pink (3%), silver (3%) and a few were green (0.3%) and violet (0.3%) (Fig. 2c). The color of microplastic in clam tissue degraded by hydrogen peroxide and potassium hydroxide found that the white, black, transparent, silver, brown, pink, and violet differed statistically ($\chi^2 = 19.057$, $df = 5$, $p = 0.002$; $\chi^2 = 13.043$, $df = 5$, $p = 0.023$; $\chi^2 = 29.728$, $df = 5$, $p = 0.000$; $\chi^2 = 11.756$, $df = 5$, $p = 0.038$; $\chi^2 = 23.777$, $df = 5$, $p = 0.000$; $\chi^2 = 24.170$, $df = 5$, $p = 0.000$; $\chi^2 = 15.256$, $df = 5$, $p = 0.009$,) whereas the light blue, navy blue, green, yellow and red, did not differ statistically ($\chi^2 = 5.545$, $df = 5$, $p = 0.353$; $\chi^2 = 8.629$, $df = 5$, $p = 0.125$; $\chi^2 = 10.084$, $df = 5$, $p = 0.073$; $\chi^2 = 7.098$, $df = 5$, $p = 0.213$; $\chi^2 = 2.807$, $df = 5$, $p = 0.730$) (Fig. 5).

The primary source determines the color of microplastics; however, they can be altered by UV radiation, weathering and microbial deterioration (Zhang et al., 2020). The colors of microplastics found in this study are similar to those found in earlier studies, where white, black and blue are the most prevalent in fishing sector equipment such as fishing nets and ground ropes (Martin et al., 2017). According to previous research, these marine organisms prefer dark-colored microplastics

because they are attracted to their color resemblance to their natural prey or to their prey, increasing the risk of ingestion (de Sá et al., 2015; Duncan et al., 2019; Ory et al., 2017).




Fig. 5 Mean concentrations of microplastic colors found in *Paratapes undulatus* after digesting clam tissue with hydrogen peroxide and potassium hydroxide

2. Microplastics' chemical composition

From the 1,001 suspected MP subsamples, FTIR analysis was employed to validate a representative number of samples (9.89%, 99 plastic items) (Table 2). The following polymer types were identified: cellulose acetate butyrate (CAB), polyethylene terephthalate (PET), bis (2-ethylhexyl) phthalate (DEHP), polypropylene (PP), cellulose acetate (CA), polyvinyl acetate (PVAc), poly (methyl methacrylate) (PMMA) and ethyl cellulose (EC) (Table 2, Fig. 6).

CAB (31.31%) was the most prevalent polymer type in the clams studied, followed by PET (9.09%). CAB is a widely used adhesive and additive in coating processes for a wide range of substrates, including plastics, textiles, metals and wood. PET is commonly used in the production of fabrics, ropes, plastic bottles, plastic bags and food containers (Qiu et al., 2015; Liu et al., 2020). According to the findings, anthropogenic trash may be a significant source of microplastics in the area where clams are harvested. PET is denser than water; it is more likely to sink and, hence, be ingested by benthic creatures. This is consistent with our results, which show that clams ingest PET.

High-molecular-weight phthalates, such as bis (2-ethylhexyl) phthalate (DEHP), are utilized in various flexible PVC daily-use products, including food packaging and home furnishings. This polymer is also found in mussels (*Mytilus galloprovincialis*) (Rios-Fuster et al., 2022).

Table 2 Polymer of microplastics identified by FTIR

Description	Number	Percentage (%)
Total particle measured (random selection)	99	100 ^a
Total polymer identified	58	58.58 ^b
Cellulose acetate butyrate (CAB)	31	31.31 ^c
Polyethylene terephthalate (PET)	9	9.09 ^c
Bis (2-ethylhexyl) phthalate (DEHP)	7	7.07 ^c
Polypropylene (PP)	5	5.05 ^c
Cellulose acetate (CA)	3	3.03 ^c
Polyvinyl acetate (PVAc)	1	1.01 ^c
Poly (methyl methacrylate) (PMMA)	1	1.01 ^c
Ethyl cellulose (EC)	1	1.01 ^c
Total non-plastic particle	41	41.42

Remark: ^aPercentage of analyzed MP particles

^bPercentage of polymers in analyzed MP particles

^cPercentage of MP polymer type

PMMA, sometimes known as acrylic glass, is a popular choice in the automotive and maritime transportation industries due to its more effective impact and ultraviolet resistance. The Gulf of Thailand, where the pier is located, is well-known for being one of Thailand's busiest shipping lanes due to its sea-based industries, such as shipping, which can obtain an abundance of plastics from commercial fisheries areas due to normal wear and tear of fishing gear, as well as commercial shipping and offshore industries derived from plastic abrasive powder and paint flakes. Moreover, as a result of nearby commercial and industrial expansion, rising urbanization has resulted in microplastic contamination in the Gulf of Thailand. In the presence of oxygen, PMMA can undergo thermal oxidative

deterioration, resulting in decreased stiffness and tensile strength, causing it to become breakable and eventually begin to develop a crack (Manoukian et al., 2019; Odli et al., 2020). Denser PMMA sinks to the seafloor and is ingested by filter feeders.

According to FTIR, 41% of the suspected microplastic particles in this analysis were non-microplastic particles. These microplastics have the potential to be confused with organic matter and other anthropogenic particles. Algae and cellulose fibers are sometimes misidentified as plastics because they break readily and contain a variety of cell layouts. In some cases, chitinous compounds that are glossy and potentially similar to plastics are not completely removed following filtration (Lusher et al., 2020). Aside from that, the centers of spherical organic particles can resemble plastic films, which can deceive our eyes. At times, salt and sand fragments may resemble plastic. Natural polymers such as wool, linen and cotton can also be confused for microplastics during microscopic analysis due to air microfiber pollution.

3. Microplastic extraction methods

Microplastics in marine species are detected by microscopically analyzing tissues or gastrointestinal contents directly (Lusher et al., 2013) or after extraction by a digestion treatment (Miller et al., 2017). Currently, a variety of digestion methods, such as acid digestion (Van Cauwenberghe & Janssen, 2014; Vandermeersch et al., 2015), KOH digestion (Foekema et al., 2013),




Fig. 6 FT-IR spectra of the representative microplastic found in clam tissue samples. Possible types are identified according to the peak position of the spectra. (A) cellulose acetate butyrate; (B) polyethylene terephthalate; (C) bis (2-ethylhexyl) phthalate; (D) polypropylene; (E) cellulose acetate; (F) polyvinyl acetate; (G) poly (methyl methacrylate); (H) ethyl cellulose

hypochlorite digestion (Collard et al., 2015), hydrogen peroxide digestion (Mathalon & Hill, 2014) and protease digestion (Cole et al., 2014), are being used to break down organic matter and make it easier to detect plastic particles. The most effective methods are those that break down tissues without damaging microplastics. In the current study, the detection of microplastics in the contents of clam tissue was made achievable by hydrogen peroxide digestion. Several studies have recently used this digesting approach for discovering microplastics in fish and mussels (Li et al., 2016; Avio et al., 2017; Güven et al., 2017). According to Tsangaris et al. (2015), preliminary testing utilizing hydrogen peroxide on virgin microplastics in powder form (PE, PP, PS and PET) revealed no changes in the appearance of plastic particles. PE, PP, PVC, PS and PET particles in the 0.3–1 mm size class had average recovery rates of 86%, which is within the range of recovery rates reported for microplastics following tissue sample digestion with hydrogen peroxide (70–95%) (Miller et al., 2017).

Although the gastrointestinal tracts of fish are frequently examined for microplastics, the entire body of small invertebrates, including mussels, shrimp and lugworms, is utilized to extract microplastics (Devriese et al., 2015; Li et al., 2015; Van Cauwenberghe et al., 2015; Phuong et al., 2018). In this research, whole clam tissue showed a larger number of microplastics per gram of tissue (4.88 ± 1.65 items g^{-1}) than whole mussel tissues of *Paratapes undulatus* (2.17 ± 0.43 to 2.38 ± 1.28 items g^{-1}) collected in Vietnam (Tran-Nguyen et al., 2023). This method can potentially detect microplastics, especially if the number of microplastics is low. This approach is also well-suited for hydrogen peroxide digestion because the time required for digestion is mostly determined by the amount of tissue to be digested (Li et al., 2015). Figure 7 depicts the organic matter-containing residue of clam tissue degraded by potassium hydroxide.




Fig. 7 Microplastic items (A-F) found in *Paratapes undulatus* after digesting clam tissue with 30% hydrogen peroxide and 10% potassium hydroxide (a-f)

Conclusion

The presence of microplastics in clams from the fresh market, where the clams were harvested from the Gulf of Thailand, indicates the ubiquitous presence of microplastics in Thai and worldwide biota. The weights of clam tissue and microplastic differed statistically significantly. The two most prevalent polymer types of microplastics identified in *P. undulatus* were polyethylene terephthalate and cellulose acetate butyrate. The quantity of microplastic discovered was consistent with earlier studies. The microplastics found in clam tissue were mostly fragments with a high percentage of sizes ranging from less than 100 µm to 200 µm, which is similar to the microplastic properties found in estuary waters. According to our results, approximately 1,011 items of microplastic per kilogram per person may enter Thai consumers' bodies from clam consumption, raising concerns about food safety and human health. It is suggested that more research be undertaken to better understand the key factors influencing microplastic abundance in clam species in order to provide effective solutions for reducing microplastic accumulation within clams.

Acknowledgements

This research is funded by Kasetsart University through the Biological Science Program in collaboration with the Faculty of Liberal Arts and Science.

References

- Avio, C.G., Gorbi, S., & Regoli, F. (2015). Experimental development of a new protocol for extraction and characterization of microplastics in fish tissues: first observations in commercial species from Adriatic Sea. *Marine Environmental Research*, 111, 18–26.
- Avio, C.G., Cardelli, L.R., Gorbi, S., Pellegrini, D., & Regoli, F. (2017). Microplastics pollution after the removal of the Costa Concordia wreck: First evidences from a biomonitoring case study. *Environmental Pollution*, 227, 207–214.
- Bayo, J., Rojo, D., & Olmos, S. (2019). Abundance, morphology and chemical composition of microplastics in sand and sediments from a protected coastal area: The Mar Menor Lagoon (SE Spain). *Environmental Pollution*, 252, 1357–1366.
- Ehlers, S.M., Manz, W., & Koop, J.H. (2019). Microplastics of different characteristics are incorporated into the larval cases of the freshwater caddisfly *Lepidostoma basale*. *Aquatic Biology*, 28, 67–77.
- Catarino, A.I., Thompson, R., Sanderson, W., & Henry, T.B. (2017). Development and optimization of a standard method for extraction of microplastics in mussels by enzyme digestion of soft tissues. *Environmental Toxicology and Chemistry*, 36, 947–951.
- Cho, Y., Shim, W.J., Jang, M., Han, G.M., & Hong, S.H. (2019). Abundance and characteristics of microplastics in market bivalves from South Korea. *Environmental Pollution*, 245, 1107–1116.
- Collard, F., Gilbert, B., Eppe, G., Parmentier, E., & Das, K. (2015). Detection of anthropogenic particles in fish stomachs: an isolation method adapted to identification by Raman spectroscopy. *Archives of Environmental Contamination and Toxicology*, 69, 331–339.
- Cole, M., Lindeque, P., Halsband, C., & Galloway, T.S. (2011). Microplastics as contaminants in the marine environment: a review. *Marine Pollution Bulletin*, 62, 2588–2597.
- Cole, M., Webb, H., Lindeque, P.K., Fileman, E.S., Halsband, C., & Galloway, T.S. (2014). Isolation of microplastics in biota-rich seawater samples and marine organisms. *Scientific Reports*, 4, 4528.
- Davidson, K., & Dudas, S.E. (2016). Microplastic ingestion by wild and cultured Manila clams (*Venerupis philippinarum*) from Baynes Sound, British Columbia. *Archives of Environmental Contamination and Toxicology*, 71, 147–156.
- de Sá, L.C., Luis, L.G., & Guilhermino, L. (2015). Effects of microplastics on juveniles of the common goby (*Pomatoschistus microps*): Confusion with prey, reduction of the predatory performance and efficiency, and possible influence of developmental conditions. *Environmental Pollution*, 196, 359–362.
- Devriese, L.I., Van der Meulen, M.D., Maes, T., Bekaert, K., Paul-Pont, I., Frère, L., ... Vethaak, A.D. (2015). Microplastic contamination in brown shrimp (*Crangon crangon*, Linnaeus 1758) from coastal waters of the Southern North Sea and Channel area. *Marine Pollution Bulletin*, 98, 179–187.
- De Witte, B., Devriese, L., Bekaert, K., Hoffman, S., Vandermeersch, G., Cooreman, K., & Robbens, J. (2014). Quality assessment of the blue mussel (*Mytilus edulis*): comparison between commercial and wild types. *Marine Pollution Bulletin*, 85, 146–155.
- Duncan, E.M., Broderick, A.C., Fuller, W.J., Galloway, T.S., Godfrey, M.H., Hamann, M., ... Godley, B.J. (2019). Microplastic ingestion ubiquitous in marine turtles. *Global Change Biology*, 25(2), 744–752.
- Falahudin, D., Cordova, M.R., Sun, X., Yogaswara, D., Wulandari, I., Hindarti, D., & Arifin, Z. (2020). The first occurrence, spatial distribution and characteristics of microplastic particles in sediments from Banten Bay, Indonesia. *Science of The Total Environment*, 705, 135304.
- Foekema, E.M., De Gruijter, C., Mergia, M.T., Van Franeker, J.A., Murk, A.J., & Koelmans, A.A. (2013). Plastic in North Sea fish. *Environmental Science & Technology*, 47, 8818–8824.

- Frias, J.P., & Nash, R. (2019). Microplastics: Finding a consensus on the definition. *Marine Pollution Bulletin*, 138, 145–147.
- Gall, S.C., & Thompson, R.C. (2015). The impact of debris on marine life. *Marine Pollution Bulletin*, 92, 170–179.
- Güven, O., Gökdağ, K., Jovanović, B., & Kıdeys, A.E. (2017). Microplastic litter composition of the Turkish territorial waters of the Mediterranean Sea, and its occurrence in the gastrointestinal tract of fish. *Environmental Pollution*, 223, 286–294.
- Harding, S. (2016). *Marine debris: Understanding, preventing and mitigating the significant adverse impacts on marine and coastal biodiversity*. Montreal, QC, Canada: Secretariat of the Convention on Biological Diversity.
- Hidalgo-Ruz, V., Gutow, L., Thompson, R.C., & Thiel, M. (2012). Microplastics in the marine environment: A review of the methods used for identification and quantification. *Environmental Science and Technology*, 46, 3060–3075.
- Hitchcock, J.N., & Mitrovic, S.M. (2019). Microplastic pollution in estuaries across a gradient of human impact. *Environmental Pollution*, 247, 457–466.
- Horton, A.A., Walton, A., Spurgeon, D.J., Lahive, E., & Svendsen, C. (2017). Microplastics in freshwater and terrestrial environments: Evaluating the current understanding to identify the knowledge gaps and future research priorities. *Science of the Total Environment*, 586, 127–141.
- Imasha, H.U.E., & Babel, S. (2023). Microplastics contamination in the green mussels (*Perna viridis*) cultured for human consumption in Thailand. *Regional Studies in Marine Science*, 67, 103203.
- Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M., Andrady, A., ... Law, K.L. (2015). Plastic waste inputs from land into the ocean. *Science*, 347(6223), 768–771.
- Jin-Feng, D., Jing-Xi, L.I., Cheng-Jun, S.U.N., Chang-Fei, H.E., Jiang, F., Feng-Lei, G.A.O., & Zheng, L. (2018). Separation and identification of microplastics in digestive system of bivalves. *Chinese Journal of Analytical Chemistry*, 46, 690–697.
- Kalogerakis, N., Karkanorachaki, K., Kalogerakis, G.C., Triantafyllidi, E.I., Gotsis, A.D., Partsinevelos, P., & Fava, F. (2017). Microplastics generation: Onset of fragmentation of polyethylene films in marine environment mesocosms. *Frontiers in Marine Science*, 4, 84.
- Karlsson, T.M., Vethaak, A.D., Almroth, B.C., Ariese, F., Van Velzen, M., Hassellöv, M., & Leslie, H.A. (2017). Screening for microplastics in sediment, water, marine invertebrates and fish: Method development and microplastic accumulation. *Marine Pollution Bulletin*, 122, 403–408.
- Lehtiniemi, M., Hartikainen, S., Nääki, P., Engström-Öst, J., Koistinen, A., & Setälä, O. (2018). Size matters more than shape: Ingestion of primary and secondary microplastics by small predators. *Food Webs*, 17, e00097.
- Leslie, H.A., Brandsma, S.H., van Velzen, M.J.M., & Vethaak, A.D. (2017). Microplastics en route: Field measurements in the Dutch river delta and Amsterdam canals, wastewater treatment plants, North Sea sediments and biota. *Environmental International*, 101, 133–142.
- Li, J., Yang, D., Li, L., Jabeen, K., & Shi, H. (2015). Microplastics in commercial bivalves from China. *Environmental Pollution*, 207, 190–195.
- Li, J., Qu, X., Su, L., Zhang, W., Yang, D., Kolandhasamy, P., Li, D., & Shi, H. (2016). Microplastics in mussels along the coastal waters of China. *Environmental Pollution*, 214, 177–184.
- Li, J., Green, C., Reynolds, A., Shi, H., & Rotchell, J.M. (2018). Microplastics in mussels sampled from coastal waters and supermarkets in the United Kingdom. *Environmental Pollution*, 241, 35–44.
- Li, J., Lusher, A.L., Rotchell, J.M., Deudero, S., Turra, A., Bråte, I.L.N., Sun, C., Hossain, M.S., ... Kolandhasamy, P. (2019). Using mussel as a global bioindicator of coastal microplastic pollution. *Environmental Pollution*, 244, 522–533.
- Liu, T., Zhao, Y., Zhu, M., Liang, J., Zheng, S., & Sun, X. (2020). Seasonal variation of micro and meso-plastics in the seawater of Jiaozhou Bay, the Yellow Sea. *Marine Pollution Bulletin*, 152, 110922.
- Lusher, A.L., McHugh, M., & Thompson, R.C. (2013). Occurrence of microplastics in the gastrointestinal tract of pelagic and demersal fish from the English Channel. *Marine Pollution Bulletin*, 67, 94–99.
- Lusher, A. (2015). Microplastics in the marine environment: Distribution, interactions and effects. In M. Bergmann, L. Gutow, & M. Klages (Eds.), *Marine anthropogenic litter* (pp. 245–307). Springer Nature.
- Lusher, A.L., Bråte, I.L.N., Munno, K., Hurley, R.R., & Welden, N.A. (2020). Is it or isn't it: The importance of visual classification in microplastic characterization. *Applied Spectroscopy*, 74(9), 1139–1153.
- Manoukian, O.S., Sardashti, N., Stedman, T., Gailiunas, K., Ojha, A., Penalosa, A., ... Kumbar, S.G. (2019). *Biomaterials for tissue engineering and regenerative medicine*. Encyclopedia of Biomedical Engineering, 2019, 462–482.
- Mathalon, A., & Hill, P. (2014). Microplastic fibers in the intertidal ecosystem surrounding Halifax Harbor, Nova Scotia. *Marine Pollution Bulletin*, 81, 69–79.
- Martin, J., Lusher, A., Thompson, R.C., & Morley, A. (2017). The deposition and accumulation of microplastics in marine sediments and bottom water from the Irish continental shelf. *Scientific Reports*, 7(1), 10772.
- Miller, M.E., Kroon, F.J., & Motti, C.A. (2017). Recovering microplastics from marine samples: A review of current practices. *Marine Pollution Bulletin*, 123, 6–18.
- Murphy, C.L. (2018). *A comparison of microplastics in farmed and wild shellfish near Vancouver Island and potential implications for contaminant transfer to humans* (Doctoral dissertation). Canada: Royal Roads University.

- Naji, A., Nuri, M., & Vethaak, A.D. (2018). Microplastics contamination in molluscs from the Northern part of the Persian Gulf. *Environmental Pollution*, 235, 113–120.
- Nor, N.H.M., & Obbard, J.P. (2014). Microplastics in Singapore's coastal mangrove ecosystems. *Marine Pollution Bulletin*, 79, 278–283.
- Odli, Z.S.M., Lee, A.J., & Asari, M.A. (2020). Marine macro- and microplastic litter along the coastal area of Kuala Perlis. In N. Samat, J. Sulong, M. Pourya Asl, P. Keikhosrokiani, Y. Azam, & S.T.K. Leng (Eds.), *Innovation and Transformation in Humanities for a Sustainable Tomorrow, Vol. 89. European Proceedings of Social and Behavioural Sciences* (pp. 303-312). European Publisher.
- Ory, N.C., Sobral, P., Ferreira, J.L., & Thiel, M. (2017). Amberstripe scad *Decapterus muroadsi* (Carangidae) fish ingest blue microplastics resembling their copepod prey along the coast of Rapa Nui (Easter Island) in the South Pacific subtropical gyre. *Science of The Total Environment*, 586, 430–437.
- Pandey, B., Pathak, J., Singh, P., Kumar, R., Kumar, A., Kaushik, S., & Thakur, T.K. (2022). Microplastics in the ecosystem: An overview on detection, removal, toxicity assessment, and control release. *Water*, 15, 51.
- Phuong, N.N., Poirier, L., Pham, Q.T., Lagarde, F., & Zalouk-Vergnoux, A. (2018). Factors influencing the microplastic contamination of bivalves from the French Atlantic Coast: Location, season and/or mode of life? *Marine Pollution Bulletin*, 129, 664–674.
- Qiu, Q., Peng, J., Yu, X., Chen, F., Wang, J., & Dong, F. (2015). Occurrence of microplastics in the coastal marine environment: First observation on sediment of China. *Marine Pollution Bulletin*, 98, 274–280.
- Qu, X., Su, L., Li, H., Liang, M., & Shi, H. (2018). Assessing the relationship between the abundance and properties of microplastics in water and in mussels. *Science of The Total Environment*, 621, 679–686.
- Renzi, M., Guerranti, C., & Blašković, A. (2018). Microplastic contents from maricultured and natural mussels. *Marine Pollution Bulletin*, 131, 248–251.
- Rios-Fuster, B., Alomar, C., Capó, X., González, G.P., Martínez, R.M.G., Rojas, D.L.S., ... Deudero, S. (2022). Assessment of the impact of aquaculture facilities on transplanted mussels (*Mytilus galloprovincialis*): Integrating plasticizers and physiological analyses as a biomonitoring strategy. *Journal of Hazardous Materials*, 424, 127264.
- Rochman, C.M., Tahir, A., Williams, S.L., Baxa, D.V., Lam, R., Miller, J.T., ... Teh, S.J. (2015). Anthropogenic debris in seafood: Plastic debris and fibers from textiles in fish and bivalves sold for human consumption. *Scientific Reports*, 5, 14340.
- Setälä, O., Norkko, J., & Lehtiniemi, M. (2016). Feeding type affects microplastic ingestion in a coastal invertebrate community. *Marine Pollution Bulletin*, 102, 95–101.
- Shahul Hamid, F., Bhatti, M.S., Anuar, N., Anuar, N., Mohan, P., & Periatnamby, A. (2018). Worldwide distribution and abundance of microplastic: How dire is the situation? *Waste Management & Research*, 36, 873–897.
- Smaal, A.C., Ferreira, J.G., Grant, J., Petersen, J.K., & Strand, Ø. (2019). Goods and services of marine bivalves. Springer: Cham, Switzerland.
- Srikrajang, J., & Prommi, T. (2021). Microplastic contamination in the edible tissues of green mussels sold in the fresh markets for human consumption. *Journal of Food Health and Bioenvironmental Science*, 14(2), 47–54.
- Su, L., Cai, H., Kolandhasamy, P., Wu, C., Rochman, C.M., & Shi, H. (2018). Using the Asian clam as an indicator of microplastic pollution in freshwater ecosystems. *Environmental Pollution*, 234, 347–355.
- Su, L., Xue, Y., Li, L., Yang, D., Kolandhasamy, P., Li, D., & Shi, H. (2016). Microplastics in Taihu lake, China. *Environmental Pollution*, 216, 711–719.
- Strady, E., Dang, T.H., Dao, T.D., Dinh, H.N., Do, T.T.D., Duong, T.N., ... Chu, V.H. (2021). Baseline assessment of microplastic concentrations in marine and freshwater environments of a developing Southeast Asian Country, Viet Nam. *Marine Pollution Bulletin*, 162, 111870.
- Suaria, G., Avio, C.G., Mineo, A., Lattin, G.L., Magaldi, M.G., Belmonte, G., ... Aliani, S. 2016). The Mediterranean plastic soup: Synthetic polymers in Mediterranean surface waters. *Scientific Reports*, 6, 37551.
- Tsangaris, C., Digka, N., Athinaïou, V., Torre, M., Anastasopoulou, A., & Zeri, C. (2015). Microplastics detection in marine biota: Methodological considerations. In *Proceedings of In the Wake of Plastics International Conference* (pp. 62). Venice: Ca' Foscari University.
- Tran-Nguyen, Q.A., Nguyen, H.N.Y., Strady, E., Nguyen, Q.T., & Trinh-Dang, M. (2020). Characteristics of microplastics in shoreline sediments from a Tropical and Urbanized Beach (Da Nang, Vietnam). *Marine Pollution Bulletin*, 161, 111768.
- Tran-Nguyen, Q.A., Nguyen, T.Q., Phan, T.L.T., Vo, M.V., & Trinh-Dang, M. (2023). Abundance of microplastics in two venus clams (*Meretrix lyrata* and *Paratapes undulatus*) from estuaries in Central Vietnam. *Water*, 15, 1312.
- Van Cauwenberghe, L., Vanreusel, A., Mees, J., & Janssen, C.R. (2013). Microplastic pollution in deep-sea sediments. *Environmental Pollution*, 182, 495–499.
- Van Cauwenberghe, L., & Janssen, C.R. (2014). Microplastics in bivalves cultured for human consumption. *Environmental Pollution*, 193, 65–70.
- Van Cauwenberghe, L., Claessens, M., Vandegehuchte, M.B., & Janssen, C.R. (2015). Microplastics are taken up by mussels (*Mytilus edulis*) and lugworms (*Arenicola marina*) living in natural habitats. *Environmental Pollution*, 199, 10–17.

- Vandermeersch, G., Van Cauwenberghe, L., Janssen, C.R., Marques, A., Granby, K., Fait, G., ... Devriese, L. (2015). A critical view on microplastic quantification in aquatic organisms. *Environmental Research*, 143, 46-55.
- Ward, J.E., Rosa, M., & Shumway, S.E. (2019). Capture, ingestion, and egestion of microplastics by suspension-feeding bivalves: A 40-year history. *Anthropocene Coasts*, 2, 39–49.
- Waser, A.M. (2018). *Predation on intertidal mussels: Influence of biotic factors on the survival of epibenthic bivalve beds*. Vrije Universiteit: Amsterdam, The Netherlands.
- Woodall, L.C., Sanchez-Vidal, A., Canals, M., Paterson, G.L., Coppock, R., Sleight, V., ... Thompson, R.C. (2014). The deep sea is a major sink for microplastic debris. *Royal Society Open Science*, 1(4), 140317.
- Zhang, K., Su, J., Xiong, X., Wu, X., Wu, C., & Liu, J. (2016). Microplastic pollution of lakeshore sediments from remote lakes in Tibet plateau, China. *Environmental Pollution*, 219, 450–455.
- Zhang, F., Man, Y.B., Mo, W.Y., Man, K.Y., & Wong, M.H. (2020). Direct and indirect effects of microplastics on bivalves, with a focus on edible species: A mini-review. *Critical Reviews in Environmental Science and Technology*, 50, 2109–2143.