

Assessment of Biodiversity among Southeast Asian Amphibians and Reptiles

HAROLD K. VORIS

*Division of Amphibians and Reptiles, Field Museum of Natural History, 1400 South Lake Shore Drive,
Chicago, IL 60605, USA*

INTRODUCTION

The 2005 Biodiversity Research and Training Program meetings featured biodiversity as a theme throughout the presentations. Since Thailand is a major center of biodiversity in Southeast Asia and is renowned for the range of its latitudes, elevations, land forms, soils, geological ages, and plant and animal communities, this was particularly appropriate.

In this paper, which is based on my presentation at those meetings, I wish to explore three important aspects of assessing biodiversity:

1. Methods and considerations in sampling biodiversity, including the roles of time and space
2. Use of traditional and modern methods to recognize species, the units of biodiversity
3. Ways of estimating the value of biodiversity

METHODS AND CONSIDERATIONS IN SAMPLING BIODIVERSITY

My work focuses on amphibians and reptiles, thus my examples are drawn from studies done on these groups. However, the issues raised, as I hope to make clear, are generally

applicable to the study of other groups and the assessment of their biodiversity.

Valid estimates of the biodiversity of amphibians and reptiles are highly dependent on the sampling methodology used. Although there is an extensive literature on the subject (see Heyer et al., 1994), sampling remains the single weakest link in the process of estimating biodiversity because it takes considerable effort to plan it thoughtfully and hard work to execute it well.

As discussed in detail by Inger and Voris (1993), Voris and Inger (1995), and summarized by Inger (2003), effective sampling surveys of amphibians and reptiles must include the following:

- Use of a variety of techniques to access the breadth of the microhabitats within each locality;
- Study of multiple localities to detect spatial diversity;
- Taking of multiple samples through time to detect seasonal variation and population oscillations.

To be specific, this means that effective surveys of amphibians and reptiles in tropical rain forests must include sampling along transects during both day and night, both in the forest and along streams, since very different species lists result from sampling these different areas at different times and in different dimensions. Microhabitats that must be sampled include tree buttresses, leaf litter, and rotting logs on the forest floor, as well as shrubs, tree trunks and vines in the vertical dimension.

*Corresponding author:

E-mail: hvoris@fieldmuseum.org




FIGURE 1. Graph showing the numbers of all frogs observed on 36 nights on the same stream (the Serbong) in Sarawak (Borneo) over a 12-month period in 1962-1963. (Inger, 2003)




FIGURE 2. Graph showing the variation in the number of individuals of one species of frog (*Limnonectes leporinus*) observed on 36 nights on the same stream (the Serbong) in Sarawak (Borneo) over a 12-month period in 1962-1963. (Inger, 2003)

Inger and Voris (1995) have documented that even streams within a few kilometers of one another that appear to be similar in topography

and microhabitat composition, have different assemblages of species. These results clearly illustrate that if sampling is done at only one or

two localities within an area, the species composition and/or relative abundance in the area will be underestimated.

Perhaps the most neglected aspect of sampling amphibians and reptiles in the tropics is the need to sample through time. The perceived lack of seasons in tropical rainforests has led to the belief that it is redundant to sample through time, but this is not the case. For example, in the area of Nanga Tekalit, in the lowland tropical rainforests of Sarawak Borneo, temperatures and rainfall are relatively constant throughout the year and there is no marked annual dry season. Yet, when Inger, in 1962, conducted a continuous full year of sampling on several hill forest streams in this setting, the resulting data clearly confirm the following assertions regarding sampling of amphibians in seemingly aseasonal tropical rainforest (Inger, 1969).

First, the level of overall activity of frogs along a stream (the Serbong) varies from night to night (Fig. 1). Thus, several nights of collecting are required to insure that samples are obtained on nights when frogs are active.

Second, different species of frogs differ in terms of their activity patterns. For example, Figure 2 illustrates the activity pattern for one species, *Limnonectes leporinus*, on the Serbong. Note that this species was observed on every night over 36 nights of collecting, although not in the same numbers.

The activity pattern for a second species, *Pedostibes hosii*, taken from the same stream, the Serbong, is shown in Figure 3. It is clear that there are many periods when this species is not active, sometimes for as long as eight consecutive nights. However on a few nights it was very active. Thus we see the necessity of making observations over a longer period of time on a single stream to insure that we detect those species that have sporadic activity patterns.

These examples illustrate and support the assertion that sampling of anurans in tropical rainforests is not a simple task but one that

requires planning and effort over a period of time.

RECOGNIZING BIODIVERSITY

Extensive sampling over multiple localities through time and space, as described above, is the crucial first step in being able to recognize individual species, the units of biodiversity.

For over 300 years, the use of traditional methods which emphasize the external morphology of amphibians and reptiles has resulted in a steady increase in the total number of species (Kohler et al., 2005; and Sodhi et al., 2004). However, in only the last 40 years, the rate of discovery of new species has escalated due to the application of new techniques and use of additional types of data obtained from behavior, ecology, karyotypes, and various molecular analyses.

Kohler et al. (2005) have carefully documented these trends for amphibians, beginning in 1758. Additional support for the strong up tick in species numbers is given in recent papers by Hanken (1999) and Bossuyt et al. (2004), which emphasize the use of molecular data to assist in the recognition of species.

Studies which I and my colleagues, Daryl Karns of Hanover College, and his students, Heather Andrews, Elizabeth Phipps, and Jerry Suddeth, have conducted on semi-aquatic snakes from within and near the Khorat Basin in northeastern Thailand, also support the notion that there are many more new snake species than previously thought.

The Khorat Basin is a prominent geographic feature of Indochina, potentially important in the ecology and evolution of semi-aquatic snake communities. We were originally interested in how different populations within known species (*Enhydrys enhydrys* and *E. plumbea*), might differ at two different but nearby geographical sites, within and outside of the Khorat Basin. We began our work using the existing definitions of species which were based on traditional morphology.

We compared community structure and population characteristics of semi-aquatic snakes (focusing on the homalopsine snakes) from a total of 11 localities located in the Khorat Basin, on the mountainous rim of the Khorat Basin, and at lower elevation sites located outside the Basin. We found that species richness (that is, numbers of species) of semi-aquatic snake assemblages

was comparable in and outside of the Khorat Basin (Table 1), and there was a high degree of species overlap between assemblages found in and outside the Basin. However, species diversity was higher in the Khorat Basin (evenness = 0.694), and *Enhydrys enhydrys* was the strong dominant at sites located outside of the Basin (evenness = 0.334).

TABLE 1. Summary of snakes collected in, on the rim of, and outside the Khorat Basin in central Thailand, June-July, 2003-04. Species are listed in overall order of abundance in each category. Numbers in parentheses under locality names indicate the number of collecting days at that site. (Taken from Karns et al., 2005)

Species	In Basin		Basin Rim			Outside Basin		Total Number of Snakes
	Khon Kaen (6)	Ban Badan (26)	Forestry Station (19)	Thai Sammakhi (19)	Bu Phram (19)	Kabin Buri (12)	Prachan Takham (4)	
Homalopsidae								
<i>Enhydris enhydris</i>	64	34	0	0	0	290	28	416
<i>Enhydris subtaeniata</i>	57	15	0	0	0	0	0	72
<i>Enhydris plumbea</i>	3	30	0	0	0	14	0	47
<i>Homalopsis buccata</i>	1	11	0	0	0	12	0	24
<i>Enhydris bocourti</i>	0	1	0	0	0	2	0	3
<i>Erpeton tentaculatum</i>	0	0	0	0	0	0	1	1
Other aquatic snakes								
<i>Xenochrophis piscator</i>	6	38	7	3	5	15	1	75
<i>Cylindrophis ruffus</i>	1	4	0	0	0	10	1	16
<i>Xenopeltis unicolor</i>	0	3	1	0	0	1	2	7
Terrestrial snakes								
<i>Oligodon taeniatus</i>	0	1	1	0	0	0	0	2
<i>Ptyas korros</i>	0	1	0	1	0	0	0	2
<i>Pareas margoritaphorus</i>	0	0	0	0	1	0	0	1
<i>Bungarus candidus</i>	0	0	1	0	0	0	0	1
<i>Elaphe flaviolineata</i>	0	0	0	0	1	0	0	1
Total Snakes	132	138	10	4	7	344	33	668
No. Snakes/Day	22.0	5.3	0.2	0.1	0.1	28.6	8.3	6.4
Species Richness	6	10	4	2	2	7	5	14
Species Evenness	0.566	0.701	0.678	0.512	0.725	0.340	0.284	

Table 1 shows that homalopsine snakes were absent from the Khorat Basin rim sites, and they were over twice as abundant at sites located outside of the Basin compared to sites inside it. We found size sexual dimorphism for *E. enhydris* outside of the Khorat Basin, but not inside it. Female snakes (*E. enhydris* and *E. plumbea*) from outside the Khorat Basin were larger and heavier compared to populations in the Basin. Historical, biogeographical, biotic, and abiotic factors that may contribute to these observed differences are discussed in Karns et al., 2005.

After collecting a total of 668 specimens throughout the course of these studies, we found some slight morphological differences between populations, and we also observed geographical differences in the distribution of the species in and out of the Basin. However,

by themselves these were insufficient to justify redefinitions of the species. It was when we conducted molecular analyses (not yet published) that significant differences among the populations were revealed, indicating that there are probably more than the initial two species we thought we were studying. Further support for the multiplication of species among these semi-aquatic snakes is provided in several recent papers describing new species in other parts of the range of this small family (Murphy & Voris, 2005; Murphy et al., 2005).

These kinds of studies and results clearly increase our estimates of the species richness and diversity present, and are an example of why the rate of discovery of new species of amphibians and reptiles has escalated in recent years.




FIGURE 3. Graph showing the variation in the number of individuals of one species of frog (*Pedostibes hosii*) observed on 36 nights on the same stream (the Serbong) in Sarawak (Borneo) over a 12-month period in 1962-1963. (Inger, 2003)

ESTIMATING THE VALUE OF BIODIVERSITY

Such careful, extensive surveys as these document the existence of even more species than previously thought, a richness of biodiversity way beyond our previous expectations. Why is this important? Why should we care? What is the value of having biodiversity and making sure it is maintained? These questions have been answered in a variety of ways

From a biological point of view, the stability of ecosystems that results from the complexity of the food chains is a very important reason to value and preserve biodiversity.

There are definitely commercial reasons to value biodiversity as well, such as maintaining sustainable harvests for the various purposes of the wildlife trade.

There is an important medical value to sustaining biodiversity both in terms of preserving sources of traditional medicines, and the potential

for discovery of sources of new drugs through bio-prospecting.

There are also significant aesthetic, historic, and recreational reasons for maintaining natural areas with their full complements of biodiversity.

But there is also another factor that is often overlooked, which we might call “the cost of construction or creation.” This is based on what was required to produce these great, complex natural systems.

Two primary elements were involved: vast amounts of space (landscape), and vast amounts of time. Plate tectonic events over the last hundreds of millions of years and episodes of glaciation over the past two million years have greatly influenced the earth’s surface, gradually developing a wide range of elevations and geological formations which, coupled with elements of latitude, drove climate changes and created abundant opportunities for the formation of species and the resultant high




FIGURE 4. An illustration showing one stage in the breakup of the supercontinent, Pangea. From a website (<http://www.scotese.com>) developed by Christopher Scotese, at the University of Texas at Arlington.

biodiversity that characterize complex ecosystems.

To illustrate some of these events in what is now Southeast Asia, we will look at three perspectives, through different spans of time.

From the late Permian period, 255 million years before the present, to the Middle Eocene epoch, some 50 million years before the present, the earth experienced the major breakup of the supercontinent, Pangea. Christopher Scotese, at the University of Texas at Arlington, has developed a website (<http://www.scotese.com>) which depicts, through a series of illustrations, the phenomenal movement of the earth's plates through this period of time. One example taken from his work is given in Figure 4.

Hall and Holloway (1998) have looked at plate tectonic shifts somewhat closer in time, from the early Eocene epoch, 55 million years before the present, to the present day, in the Austral-Asia region. A compact disc which

accompanies their paper gives an impressive animated portrayal of these events. Fig. 5 illustrates one view from their work.

The third perspective, given by Wallace (1881), Molengraaff & Weber (1919), Darlington (1957), Voris, (2000) and Sathiamurthy & Voris (in press, 2006), focuses on the sea level and climate changes between the period from 250,000 years ago, to the present day. Changes in sea levels over geological time have long been considered crucial to understanding the distribution of both aquatic and terrestrial organisms that we see today.

Figure 6 from Voris (2000), gives a map of present day Austral-Asia, along with a depiction of the extent of the Sunda and Sahul shelves when sea levels were 75 meters below present day levels, showing the magnitude of change in the land mass between two points in time. The publication gives numerous maps of the region at other sea levels.




FIGURE 5. The Austral-Asia region at 40 million years before the present as depicted by Hall and Holloway (1998). A compact disc that accompanies their publication contains this illustration and many others.




FIGURE 6. A map of present-day Austral-Asia showing the extent of the Sunda and Sahul shelves when sea levels were 75 meters below present-day levels. (Voris, 2000)




FIGURE 7. A map of present-day Austral-Asia showing the extent of the exposed Sunda Shelf due to lower sea levels at 21 ka years before present. (Sathiamurthy and Voris, in press, 2006)

Sathiamurthy and Voris (in press, 2006) have further refined and expanded these maps using a comprehensive data base of ocean

depths. Figure 7 is one example from their work, again showing a map of present-day Austral-Asia, along with the extent of the Sunda

Shelf when sea levels were even lower than in the above example from Voris (2000), some 21 thousand years before the present.

It is impossible to fully convey in a written paper the awesome impression made by seeing all the visual depictions shown in these studies. However, we can begin to appreciate the magnitude of time and events that have led to the earth's present configuration of geography and ecosystems, by referring to our own experiences of how the powerful forces that result from plate tectonic shifts, volcanic activity, earthquakes, and tsunamis continue to reshape our environment. We cannot help but be led to a profound appreciation of what the earth has undergone.

We are led to an inescapable conclusion: the enormous amounts of time and spatial diversity on this planet cannot be duplicated, making the biodiversity that has resulted, absolutely unique, irreplaceable, and priceless.

ACKNOWLEDGEMENTS

This work was supported by the Field Museum of Natural History, the Hanover College Biology Department, and the Rivers Institute at Hanover College. I thank Daryl Karns and John Murphy for their close collaboration and their many contributions to this work both in the field and laboratory. I thank my many colleagues and sponsors in Thailand. In particular, I thank the National Research Council of Thailand; Mr. Taksin and his staff at the Sakaerat Environmental Research Station; Jarujin Nabhitabhata, Tanya Chanard, and Yodchaiy Chuaynkern at the National Science Museum; Lawan Chanhome at the Queen Saovabha Memorial Institute, Thai Red Cross Society; Chantip Inthara in the Department of Biology at Khon Kaen University; Dr. Vachira Lheknim, at the Prince of Songkhla University; and Dr. Kumthorn Thirakhupt, Dr. Somsak Panha, and Dr. Sansareeya Wangkulangkul in the Department of Biology at Chulalongkorn University. At the field sites, assistance from Jacques Hill, David

McLeod, and Teresa Mayfield was much appreciated. I also appreciate the hard labor of the many snake collectors who assisted us in the various towns where we worked, especially Pom Mongodjng and Vera Dummai from Ban Badan and Utid Naenon from Kabin Buri. We were ably assisted in all aspects of this study by our driver translator research colleague, Prapon (Pon) Noiwan. At the Field Museum special thanks for assistance are due to Tom Anton, Jamie Ladonski, Alan Resetar, Jennifer Mui, Kayte O'Brien, and Sarah Drasner. I thank Bob Inger for permission to use illustrations from one of his papers. Thanks also go to Hanover College students Heather Andrews, Elizabeth Phipps, and Jerry Suddeth. I give special thanks to Helen Voris for her able assistance in the preparation of the B.R.T. talk and the writing and editing of this manuscript. In addition, I thank Alan Resetar for his careful review of the manuscript.

LITERATURE CITED

- Bossuyt, F., Meegaskumbura, M., Beenaerts, N., Gower, D.J., Pethiyagoda, R., Roelants, K., Mannaert, A., Wilkinson, M., Bahir, M.M., Manamendra-Arachchi, K., Ng, P.K.L., Schneider, C.J., Oommen, O.V., and Milinkovitch, M.C. 2004. Local endemism within the western Ghats-Sri Lanka biodiversity hotspot. *Science*, 306: 479-481.
- Darlington, P.J. 1957. *Zoogeography: The Geographical Distribution of Animals*. John Wiley & Sons, Inc., New York.
- Hall, R., and Holloway, J.D. (Eds.). 1998. *Biogeography and Geological Evolution of SE Asia*, p. 417. Backhuys Publishers, Leiden.
- Hanken, J. 1999. Why are there so many new amphibian species when amphibians are declining? *Tree*, 14: 7-8.
- Heyer, W.R., Donnelly, M.A., McDiarmid, R.W., Hayek, L.C., and Foster, M.S. (Eds.). 1994. *Measuring and Monitoring Biological Diversity Standard Methods for Amphibians*. Biological Diversity Handbook Series. Smithsonian Institution Press, Washington & London.
- Inger, R.F. 1969. Organization of communities of frogs along small rain forest streams in Sarawak. *Journal Animal Ecology*, 38: 123-148.

- Inger, R.F., 2003. Sampling biodiversity in Bornean frogs. *The Natural History Journal of Chulalongkorn University*, 3(1):9-15.
- Inger, R.F., and Voris, H.K. 1993. A comparison of amphibian communities through time and from place to place in Bornean forests. *Journal of Tropical Ecology*, 9: 409-433.
- Karns, D.R., Murphy, J.C., Voris, H.K., and Suddeth, J.S. 2005. Comparison of semi-aquatic snake communities associated with the Khorat Basin, Thailand. *The Natural History Journal of Chulalongkorn University*, 5: 73-90.
- Kohler, J., Vieites, D.R., Bonett, R.M., Garcia, F.H., Glaw, F., Steinke, D., and Vences, M. 2005. New amphibians and global conservation: a boost in species discoveries in a highly endangered vertebrate group. *BioScience*, 55: 693-696.
- Molengraaff, G.A.F., and Weber, M. 1919. Het verband tusschen den plistoceenen ijstijd en het ontstaan der Soenda-zee (Java- en Zuid-Chineesche Zee) en de invloed daarvan op de verspreiding der koraalriffen en op de land-en zoetwater-fauna. *Verslag Van de Gewone Vergaderingen der Wis-En Natuurkundige Afdeeling*, 28: 497-544.
- Murphy, J.C., and Voris, H.K. 2005. A new Thai *Enhydryis* (Serpentes: Colubridae: Homalopsinae). *Raffles Bulletin of Zoology*, 53: 143-147.
- Murphy, J.C., Voris, H.K., and Auliya, M. 2005. A new species of *Enhydryis* (Serpentes: Colubridae: Homalopsinae) from the Kapuas river system, West Kalimantan, Indonesia. *Raffles Bulletin of Zoology*, 53: 115-119.
- Sathiamurthy, E., and Voris, H.K. 2006. Maps of Holocene sea level transgression and submerged lakes on the Sunda Shelf. *The Natural History Journal of Chulalongkorn University*, (In press).
- Scotese, C.R., 2005. Website at <http://www.scotese.com/>, University of Texas at Arlington.
- Sodhi, N.S., Koh, L.P., Brook, B.W., and Ng, P.K.L. 2004. Southeast Asian biodiversity: an impending disaster. *Trends in Ecology and Evolution*, 19: 654-660.
- Voris, H.K. 2000. Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems, time durations. *Journal of Biogeography*, 27: 1153-1167.
- Voris, H.K., and Inger, R.F. 1995. Frog abundance along streams in Bornean forests. *Conservation Biology*, 9: 679-683.
- Wallace, A.R. 1881. *Island Life or the Phenomena and Causes of Insular Faunas and Floras Including a Revision and Attempted Solution of the Problem of Geological Climates*. Macmillan, London.

Received: 19 April 2006

Accepted: 3 May 2006